

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 452 530 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
01.09.2004 Bulletin 2004/36

(21) Application number: 02783753.3

(22) Date of filing: 03.12.2002

(51) Int Cl.7: **C07D 277/42**, C07D 417/04,
C07D 417/12, A61K 31/426,
A61K 31/427, A61K 31/454,
A61K 31/496, A61K 31/5377,
A61K 31/4439, A61P 3/06,
A61P 3/10, A61P 43/00

(86) International application number:
PCT/JP2002/012673

(87) International publication number:
WO 2003/048140 (12.06.2003 Gazette 2003/24)

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
IE IT LI LU MC NL PT SE SI SK TR
Designated Extension States:
AL LT LV MK RO

(30) Priority: 03.12.2001 JP 2001368567

(71) Applicant: **Japan Tobacco Inc.**
Tokyo 105-8422 (JP)


(72) Inventors:
• **INABA, T.**, Japan Tobacco Inc.
Takatsuki-shi, Osaka 569-1125 (JP)

- **IKEMOTO, Tomoyuki**, Japan Tobacco Inc.
Takatsuki-shi, Osaka 569-1125 (JP)
- **SAKATA, Shohei.**, Japan Tobacco Inc.
Takatsuki-shi, Osaka 569-1125 (JP)
- **MAEGAWA, Hiroshi**
Kusatsu-shi, Shiga 525-0072 (JP)
- **KASHIWAGI, Atsunori**
Ohtsu-shi, Shiga 520-2264 (JP)

(74) Representative:
von Kreisler, Alek, Dipl.-Chem. et al
Deichmannhaus am Dom,
Postfach 10 22 41
50462 Köln (DE)

(54) **AZOLE COMPOUND AND MEDICINAL USE THEREOF**

(57) The present invention relates to an azole compound represented by the formula [I]



wherein W is S or O; R is -COOR⁷, -X¹-A¹-COOR⁷ (R⁷ is H, alkyl) or tetrazolyl; R¹, R², R³ and R⁴ are H and the like; A is -(CH₂)_m-X- (X is -N(R⁸)-, -C(R⁹)(R¹⁰)-, -CO- or -CO-N(R⁸)-); B is aryl or aromatic heterocyclic group; R⁵ is H and the like; R⁶ is -(Y)_{s1}-(A²)_s-Z (Y is -O-, -S(O)_t-, -N(R¹³)-, -N(R¹⁴)-CO-, -N(R¹⁴)-SO₂-, -SO₂-N(R¹⁴)- and the like, A² is alkylene, and Z is cycloalkyl, aryl, aromatic heterocyclic group, indanyl, piperazinyl, a prodrug thereof or a pharmaceutically acceptable salt thereof. The compound [I] of the present invention has a protein tyrosine phosphatase 1B inhibitory activity, and is useful as a therapeutic agent for diabetes, a therapeutic drug for diabetic complications or a therapeutic drug for hyperlipidemia.

BEST AVAILABLE COPY

EP 1 452 530 A1

Description**Technical Field**

5 **[0001]** The present invention relates to a novel azole compound, more particularly, the present invention relates to an azole compound or a pharmaceutically acceptable salt thereof having a protein tyrosine phosphatase 1B (PTP1B) inhibitory activity, and a pharmaceutical composition containing the same.

Background Art

10

[0002] Diabetes causes various metabolic abnormalities including a chronic hyperglycemia state as a main characteristic, which shows various symptoms based on mouth dryness, polydipsia, polyuria, decrease in body weight and the like, based on hyperglycemia. It is known that, when such hyperglycemia state is maintained for a long time, various complications such as retinopathy, nephropathy, neuropathy, cardiac infarction and cerebral infarction based on arteriosclerosis, and the like are developed.

15

[0003] Diabetes is roughly divided into four types of type I diabetes (IDDM; Insulin Dependent Diabetes Mellitus), wherein pancreatic beta cells are damaged or destroyed to cause absolute insulin deficiency, type II diabetes (NIDDM; Non-Insulin Dependent Diabetes Mellitus) wherein relative insulin deficiency is developed due to insulin resistance and lower insulin secretion, specific diabetes which secondarily results from genetic abnormalities, other diseases and the like, and gestational diabetes. Some of those diagnosed with type II diabetes at the time of the onset may gradually lose their ability to secrete insulin with the progression of the disease and result in type I diabetes.

20

[0004] Considering the saccharometabolism of living organisms, energy sources and materials to be the constituent components of living organisms are taken into the body intermittently; for example, the brain keeps on consuming glucose. Under such situation, the blood glucose level is maintained almost constantly, and what enables such control of the blood glucose level includes hormones involved in control of the blood glucose level, metabolism in organs, interaction of exchanging sugar and the like between organs. Of such hormones, particularly the action of insulin involved in the control of the blood glucose level is important, and its disorder, namely, insulin resistance and lower secretion of insulin are considered to be deeply involved in diabetes.

25

[0005] Insulin is secreted from pancreatic beta cells, binds with an insulin receptor present on the membrane surface of the skeletal muscle cell and adipocyte, which are its target cells, after which tyrosine residue in the intracellular domain is self-phosphorylated. Then, tyrosine residues such as insulin receptor substrate (IRS), APS (adapter protein containing PH and SH2 domain) and the like are phosphorylated and PI₃ kinase -Akt pathway is activated, which causes translocation of glucose transporter to cell membrane, where glucose uptake occurs to lower the blood glucose level. On the other hand, tyrosine phosphatase that causes tyrosine dephosphorylation to negatively control intracellular signal transduction by insulin is also present, and suppresses activation thereof. In this way, while tyrosine phosphorylation plays a key role in the insulin action, considering that tyrosine phosphorylation depends on the activity balance between tyrosine kinase (phosphorylation enzyme) and tyrosine phosphatase (dephosphorylation enzyme), tyrosine phosphatase is presumed to have function to significantly control insulin signal transduction directly together with tyrosine kinase.

35

[0006] At present, tyrosine phosphatase forms a large gene family and more than 70 some kinds of isozyme have been reported. Of such isozymes, protein tyrosine phosphatase 1B (PTP1B) is considered to be a phosphatase specific to insulin signal transduction. Particularly, given the reports on increased gene expression of PTP1B by high glucose culture, and shift of intracellular localization thereof, which decreases insulin receptor and IRS-1 tyrosine phosphorylation and induces insulin resistance (J. Biol. Chem., 270: 7724-7730, 1995; J. Biochem. (Tokyo), 123: 813-820, 1998); and introduction of wild-type PTP1B prevents translocation of glucose transporter GLUT4, resulting in ineffectiveness in a phosphatase activity deficient mutant, and recently on enhanced insulin sensitivity of PTP1B knockout mouse to be obesity resistant to high-fat diet (Science, 283: 1544-1548, 1999), this enzyme is suggested to be one possible target of insulin resistance improvement. In fact, an insulin resistance improvement effect of vanadic acid long known as a tyrosine phosphatase inhibitor has been acknowledged in animal test and the like.

45

[0007] Accordingly, such tyrosine phosphatase, particularly a drug that suppresses and/or inhibits abnormal activation of PTP1B can be a new type of therapeutic agent for diabetes, which enhances insulin sensitivity, insulin resistance and/or glucose resistance, and restores normal insulin intracellular signal transduction. In addition, application to a therapeutic drug for various diseases such as obesity, neurodegenerative disease and the like is also expected.

50

[0008] Recently, various reports have been documented on compounds aiming at treatment of diseases such as diabetes and the like, by inhibiting protein tyrosine phosphatase.

55

[0009] For example, WO 00/17211 discloses a phosphonic acid derivative having a PTP1B inhibitory action. However, this publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof.

[0010] JP-11-508919A (US 5,770,620) discloses an arylacrylic acid derivative useful as a protein tyrosine phosphatase inhibitor. However, this publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof.

[0011] WO 98/27092 (US 6,080,772) discloses a thiazole compound having a protein tyrosine phosphatase inhibitory action. However, this publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof.


[0012] WO 99/58522 discloses a naphtho[2,3-B]heteroal-4-yl derivative, WO 99/58511 discloses an oxa/thiazole-aryl-carboxylic acid derivative, WO 99/58521 and US 6,110,962 disclose 11-aryl-benzo[B]naphtho[2,3-D]furan and 11-aryl-benzo[B]naphtho[2,3-D]thiophene derivatives, WO 99/58518 discloses a biphenyl-oxo-acetic acid derivative, WO 99/61419 discloses a 2,3,5-substituted biphenyl derivative, WO 99/58520 discloses a biphenyl-sulfonyl-aryl-carboxylic acid derivative, WO 99/61435 discloses benzothiophene, benzofuran and indole derivatives, US 6,103,708 discloses furan, benzofuran and thiophene derivatives, US 6,110,963 discloses an aryl-oxo-acetic acid derivative, US 6,001,867 discloses a 1-aryl-dibenzothiophene derivative, US 6,057,316 discloses a 4-aryl-1-oxa-9-thia-cyclopenta[B]fluorene derivative, US 6,063,815 discloses a benzophenone derivative, as each having a protein tyrosine phosphatase inhibitory action. However, these publications do not disclose compounds having a structure as that of the compound of the present invention, not to mention a description suggestive thereof.

[0013] As compounds having a thiazole or oxazole structure, the following have been reported.

[0014] WO 00/45635 discloses a 2-substituted thiazole derivative. However, the compound of this publication has a carbamoyl group at the terminal of the substituent at the 2-position of the thiazole ring and this publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof. In addition, the compound of this publication is useful as an antimicrobial agent or an analgesic, and the publication does not disclose its usefulness as a PTP1B inhibitor, not to mention a description suggestive thereof.

[0015] JP-2000-504039A describes a 2-anilino-4-phenylthiazole derivative. However, the compound of this publication has an anilino group substituted by a hydroxyl group or a carboxyl group at the 2-position of a thiazole ring, a phenyl group at the 4-position, and a substituent at the 2-position of the 4-position phenyl group, and this publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof. In addition, the compound of this publication is useful as a CRF (corticotropin releasing factor) antagonist, and the publication does not disclose its usefulness as a PTP1B inhibitor, not to mention a description suggestive thereof.


[0016] JP-A-4-154773 describes a thiazole derivative represented by the formula




wherein R^1 and R^2 are the same or different and each is hydrogen atom, halogen atom, lower alkyl group, phenyl group, substituted phenyl group, pyridyl group or substituted pyridyl group, R^3 is hydroxyl group, lower alkoxy group or $-N(R^5)(R^6)$ wherein R^5 and R^6 are the same or different and each is hydrogen atom or lower alkyl group, R^4 is hydrogen atom or lower alkyl group, and X is amino group, amide group, carbonyl group, alkylene group, oxygen atom or sulfur atom. However, this publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof. In addition, the compound of this publication is useful as an antiinflammatory agent, and the publication does not disclose its usefulness as a PTP1B inhibitor, not to mention a description suggestive thereof.

[0017] WO 94/08982 describes a 4-phenylthiazole derivative. However, the compound of this publication has phenyl group at the 4-position of a thiazole ring, and a substituent such as halogen and the like at the 2-position of the 4-position phenyl group. This publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof. In addition, the compound of this publication is useful as a pest control agent, and the publication does not disclose its usefulness as a PTP1B inhibitor, not to mention a description suggestive thereof.

[0018] WO 02/39997 describes compounds represented by the formula




wherein R^6 is hydroxyl group or protected prodrug moiety, R^7 is hydrogen atom, carboxy group, arylaminocarbonyl group, aroyl group, aryl group, alkylaminocarbonyl group, aminocarbonyl group, alkenylaminocarboxy group, hydroxyl group, alkoxy group, ether, thiol, amino group-containing heterocyclic group or protected prodrug moiety, R^8 is hydrogen atom or alkyl group that may be bonded with D to form a ring, R^9 is lower alkyl group or hydrogen atom, Q is bond, oxygen atom, sulfur atom, CR^3OH , CR^3SH , $CR^3NR^{3a}R^{3b}$, NR^3 , $(CR^3R^{3a})_n$, $O(CR^3R^{3b})_n$ or $(CR^3R^{3a})_nO(CR^{3b}R^{3c})_n$ wherein n is an integer of 0 or 1 to 3, R^3 , R^{3a} , R^{3b} and R^{3c} are each independently hydrogen atom, optionally substituted straight chain, cyclic or branched chain C_{1-6} alkyl group, C_{2-6} alkenyl group, acyl group, arylalkyl group, aryloxy carbonyl group, arylaminocarbonyl group, arylalkylsulfonyl group or aryl group, G is a linking moiety, M is anchor moiety, J is bond, alkylene group, alkenylene group or alkynylene group, D is hydrogen atom, alkoxy group, amine, alkyl group, alkenyl group, alkynyl group, aryl group or heteroaryl group that may be bonded with G, M or Q to form a ring, t is 0 or 1, p is 0 or an integer of 1 to 5, and q is 0 or an integer of 1 to 3, and the formula



wherein P^4 is carboxy group, cleavable prodrug moiety, $COOP^4$, $(CH_2)_{1-4}SP^4$ or $C(O)NP^4P^4$, R^7 is hydrogen atom, carboxy group, optionally substituted lower alkyl ester, lower alkenyl ester, ester added with secondary amine substituted by lower alkyl, arylaminocarbonyl group, aroyl group, aryl group, alkylaminocarbonyl group, aminocarbonyl group, $COOR^7$, $CONR^7R^7$, hydroxyl group, ether, thiol, amino group, $(CH_2)_{1-4}SR^7$, heterocyclic group or cleavable prodrug moiety, P^4 , P^4 , R^7 and R^7 are each independently hydrogen atom, C_{1-6} alkyl group, C_{2-6} alkenyl group, C_{2-6} alkynyl group or optionally substituted aryl group, R^8 is hydrogen atom, alkyl group or covalent bond with D, R^9 is lower alkyl group or hydrogen atom, Q is bond, oxygen atom, sulfur atom, CR^3OH , CR^3SH , $CR^3NR^{3a}R^{3b}$, NR^3 , $(CR^3R^{3a})_n$, $O(CR^3R^{3b})_n$ or $(CR^3R^{3a})_nO(CR^{3b}R^{3c})_n$ wherein n is 0 or an integer of 1 to 3, R^3 , R^{3a} , R^{3b} and R^{3c} are each independently hydrogen atom, optionally substituted C_{1-6} straight chain or branched chain alkyl group, C_{2-6} straight chain or branched chain alkenyl group, aryloxy carbonyl group, arylaminocarbonyl group, arylalkylsulfonyl group, arylalkyl group, optionally substituted acyl group, aryl group or C_{3-8} ring optionally substituted by up to 4 hetero atoms, P^{2a} , P^{2b} , P^{3a} and P^{3b} are each independently hydrogen atom or optionally substituted straight chain, branched chain or cyclic C_{1-5} alkyl group, G is linking moiety, M is anchor moiety, J is bond, alkylene group, alkenylene group or alkynylene group, D is hydrogen atom, alkyl group, alkenyl group, alkynyl group or aryl group or may be bonded with G, M or Q to form a ring, t is 0 or 1, p is 0 or an integer of 1 to 5, and q is 0 or an integer of 1 to 3, wherein the anchor moiety in each formula is exemplified by thiazole group and oxazole group having, as a substituent, aryl group or heteroaryl group substituted by $-NR'R''$, $-CONR'R''$, $-S(O)_2NR'R''$, $-S(O)_{0-2}R'$, $-NR'R''$, $-O(CR'R'')_{0-2}CF_3$, $-COR'$, $-CO_2R'$ and $-OR'$ wherein R' and R'' are each independently hydrogen atom, C_{1-6} alkyl group, C_{2-6} alkenyl group, C_{2-6} alkynyl group or optionally substituted aryl group, and the linking moiety is exemplified by covalent bond and C_{1-6} alkyl group.

[0019] Moreover, a compound represented by the formula



wherein M is carbocyclic group, heterocyclic group or CONR'R'' wherein R' and R'' are each independently hydrogen atom, C₁₋₆ alkyl group, C₂₋₆ alkenyl group, C₂₋₆ alkynyl group or optionally substituted aryl group, Q is bond, oxygen atom, sulfur atom, CR³OH, CR³SH, CR³NR^{3a}R^{3b}, NR³, (CR³R^{3a})_n, O(CR³R^{3b})_n or (CR³R^{3a})_nO(CR^{3b}R^{3c})_n wherein n is 0 or an integer of 1 to 3, R³, R^{3a}, R^{3b} and R^{3c} are each independently hydrogen atom, optionally substituted branched chain, cyclic or straight chain C₁₋₆ alkyl group, C₂₋₆ alkenyl group, acyl group, arylalkyl group, aryloxy carbonyl group, arylaminocarbonyl group, arylalkylsulfonyl group or aryl group, K is independently selected sublinking moiety, L is independently selected subanchor moiety, P⁴ is hydrogen atom, carboxy group, (CH₂)₁₋₄SP⁴, cleavable prodrug moiety, COOP⁴ or CONP⁴P⁴, R⁷ is hydrogen atom, carboxy group, aroyl group, aryl group, COOR⁷, C(O)NR⁷R⁷, hydroxyl group, ether, thiol, (CH₂)₁₋₄SR⁷, heterocyclic group or cleavable prodrug moiety, P⁴, P⁴, R⁷ and R⁷ are each independently hydrogen atom, C₁₋₆ alkyl group, C₂₋₆ alkenyl group, C₂₋₆ alkynyl group or optionally substituted aryl group, n is 0 or an integer of 1 to 4, D is hydrogen atom, alkyl group, alkoxy group, alkenyl group, amine, hydroxyl group, alkynyl group, aryl group or heteroaryl group, and t is 0 or 1, is described, wherein the sublinking moiety has a covalent bond and the subanchor moiety has an optionally substituted aryl group.

[0020] However, this publication does not disclose a compound having a structure as that of the compound of the present invention, not to mention a description suggestive thereof. In addition, the compound of this publication is useful as an angiotensin converting enzyme (ACE)-2 regulator, and the publication does not disclose its usefulness as a PTP1B inhibitor, not to mention a description suggestive thereof.

Disclosure of the Invention


[0021] An object of the present invention is to provide a compound having a superior PTP1B inhibitory action and useful as a therapeutic agent for diabetes, a therapeutic agent for hyperlipidemia or a therapeutic drug of diseases such as obesity, neurodegenerative disease and the like.

[0022] Another object of the present invention is to provide a PTP1B inhibitor, a therapeutic agent for diabetes and a therapeutic agent for hyperlipidemia.

[0023] The present inventors have conducted intensive studies in an attempt to achieve the above-mentioned objects and found that an azole compound represented by the following formula [I] has a superior PTP1B inhibitory action and is useful as a PTP1B inhibitor, a therapeutic agent for diabetes and a therapeutic agent for hyperlipidemia, which resulted in the completion of the present invention.

[0024] The present invention relates to the compounds shown in the following [1] to [54] and use thereof as a pharmaceutical agent.

[1] An azole compound represented by the formula [I]



wherein

W is a sulfur atom or an oxygen atom;

R is

(1) $-\text{COOR}^7$ wherein R^7 is a hydrogen atom or a lower alkyl group),

(2) $-\text{X}^1-\text{A}^1-\text{COOR}^7$

wherein

X^1 is $-\text{O}-$, $-\text{N}(\text{R}^{15})-$ or $-\text{S}(\text{O})_p-$ wherein R^{15} is a hydrogen atom or a lower alkyl group, p is 0, 1 or 2,

A^1 is a lower alkylene group, and

R^7 is a hydrogen atom or a lower alkyl group, or

(3) a tetrazolyl group;

R^1 , R^2 , R^3 and R^4 are each independently

(1) a hydrogen atom,

(2) a halogen atom,

(3) a hydroxyl group,

(4) an optionally substituted lower cycloalkylalkyloxy group,

(5) an optionally substituted aralkyloxy group,

(6) a cyano group,

(7) a nitro group,

(8) a lower alkyl group,

(9) a lower haloalkyl group,

(10) a lower alkoxy group or

(11) a lower haloalkoxy group;

A is a group represented by $-(\text{CH}_2)_m-\text{X}-$

wherein

X is $-\text{N}(\text{R}^8)-$, $-\text{C}(\text{R}^9)(\text{R}^{10})-$, $-\text{CO}-$ or $-\text{CO}-\text{N}(\text{R}^8)-$

wherein

R^8 is a hydrogen atom, $-\text{SO}_2\text{R}^{16}$ (R^{16} is a lower alkyl group or an aryl group) or a lower alkyl group, wherein said lower alkyl group is optionally substituted by a substituent selected from the group consisting of a lower alkoxy group, an aryloxy group, $-\text{N}(\text{R}^{11})(\text{R}^{12})$ (R^{11} and R^{12} are each independently a hydrogen atom or a lower alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom),

a carboxy group, a lower cycloalkyl group and an optionally substituted aryl group, and

R^9 and R^{10} are each independently a hydrogen atom or a lower alkyl group or may form lower cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, and

m is 0 or an integer of 1 to 3;

B is an aryl group or an aromatic heterocyclic group;

R^5 is

(1) a hydrogen atom,

(2) a halogen atom,

(3) a lower alkyl group,

(4) a lower alkoxy group,

(5) a cyano group,

- (6) a nitro group,
- (7) a lower haloalkyl group or
- (8) $-S(O)_t-R^{17}$ (R^{17} is a lower alkyl group or an aryl group and t is 0, 1 or 2); and

R^6 is $-(Y)_{s1}-(A^2)_s-Z$

wherein

$s1$ and s are each independently 0 or 1,

Y is $-O-$, $-S(O)_t-$, $-N(R^{13})-$, $-N(R^{14})-CO-$, $-N(R^{14})-SO_2-$, $-SO_2-N(R^{14})-$, $-C(R^{18})(R^{19})-$ or $-CO-$

(wherein

t is 0, 1 or 2,

R^{13} is

- (1) a hydrogen atom,
- (2) a lower alkyl group

(wherein said lower alkyl group optionally substituted by a substituent selected from the group consisting of

- (a) a lower cycloalkyl group,
- (b) an optionally substituted aryl group,
- (c) an optionally substituted heterocyclic group and
- (d) a hydroxyl group),

- (3) a lower alkenyl group,
- (4) a lower alkylsulfonyl group or
- (5) a lower alkylcarbonyl group

(wherein said lower alkylcarbonyl group is optionally substituted by a hydroxyl group or a lower alkoxy group),

R^{14} is a hydrogen atom or a lower alkyl group, and

R^{18} and R^{19} are each independently a hydrogen atom or a lower alkyl group or may form lower cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom),

A^2 is a lower alkylene group optionally substituted by a lower cycloalkyl group, and

Z is

- (1) a lower cycloalkyl group
- (wherein said a lower cycloalkyl group is optionally substituted by an optionally substituted phenyl group),
- (2) an aryl group
- (wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group optionally substituted by a substituent selected from the group consisting of a lower alkyl group and a lower alkylcarbonyl group,
- (b) a lower cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a lower alkyl group,
- (c) a carboxy group,
- (d) a halogen atom,
- (e) an alkyl group,
- (f) a lower haloalkyl group,
- (g) a lower alkylamino group,
- (h) a di(lower alkyl)amino group,
- (i) a lower alkylthio group and
- (j) a lower alkoxy group),

- (3) an optionally substituted aromatic heterocyclic group,
- (4) an indanyl group or
- (5) a piperazinyl group

(wherein said piperazinyl group is optionally substituted by a substituent selected from the group consisting of

- (a) a phenyl group,
- (b) a phenyl lower alkyl group,
- (c) a benzoyl group optionally substituted by a halogen atom and
- (d) a phenyl lower alkoxy carbonyl group),

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[2] The azole compound of [1], wherein, in the formula [I], W is a sulfur atom or an oxygen atom;

R is

- (1) -COOR⁷ wherein R⁷ is a hydrogen atom or a C₁₋₄ alkyl group,
- (2) -X¹-A¹-COOR⁷
wherein
X¹ is -O-, -N(R¹⁵)- or -S(O)_p- wherein R¹⁵ is a hydrogen atom or a C₁₋₄ alkyl group and p is 0, 1 or 2,
A¹ is a C₁₋₄ alkylene group,
R⁷ is a hydrogen atom or a C₁₋₄ alkyl group or
- (3) a tetrazolyl group,

R¹, R², R³ and R⁴ are each independently,

- (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a hydroxyl group,
- (4) an optionally substituted C₃₋₇ cycloalkyl C₁₋₄ alkoxy group,
- (5) an optionally substituted aralkoxy group,
- (6) a cyano group,
- (7) a nitro group,
- (8) a C₁₋₄ alkyl group,
- (9) a C₁₋₄ haloalkyl group,
- (10) a C₁₋₄ alkoxy group or
- (11) a C₁₋₄ haloalkoxy group;

A is a group represented by -(CH₂)_m-X-

wherein

X is -N(R⁸)-, -C(R⁹)(R¹⁰)-, -CO- or -CO-N(R⁸)-

wherein

R⁸ is a hydrogen atom, -SO₂R¹⁶ (R¹⁶ is a C₁₋₆ alkyl group or an aryl group) or a C₁₋₆ alkyl group, wherein said C₁₋₆ alkyl group is optionally substituted by a substituent selected from the group consisting of a C₁₋₄ alkoxy group, an aryloxy group, -N(R¹¹)(R¹²) (R¹¹ and R¹² are each independently a hydrogen atom or a C₁₋₄ alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a C₃₋₇ cycloalkyl group and an optionally substituted aryl group,
R⁹ and R¹⁰ are each independently a hydrogen atom or a C₁₋₄ alkyl group, or may form a C₃₋₇ cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, m is 0 or an integer of 1 to 3;

B is an aryl group or an aromatic heterocyclic group;

R⁵ is

- (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a C₁₋₄ alkyl group,
- (4) a C₁₋₄ alkoxy group,
- (5) a cyano group,
- (6) a nitro group,
- (7) a C₁₋₄ haloalkyl group or
- (8) -S(O)_r-R¹⁷ (R¹⁷ is a C₁₋₆ alkyl group or an aryl group and r is 0, 1 or 2);

R⁶ is -(Y)_{s1}-(A²)_s-Z

wherein

s1 and s are each independently 0 or 1,

Y is -O-, -S(O)_t-, -N(R¹³)-, -N(R¹⁴)-CO-, -N(R¹⁴)-SO₂-, -SO₂-N(R¹⁴)-, -C(R¹⁸)(R¹⁹)- or -CO-

(wherein

t is 0, 1 or 2,

R¹³ is

(1) a hydrogen atom,

(2) a C₁₋₄ alkyl group

(wherein said C₁₋₄ alkyl group is optionally substituted by a substituent selected from the group consisting of

(a) a C₃₋₇ cycloalkyl group,

(b) an optionally substituted aryl group,

(c) an optionally substituted heterocyclic group and

(d) a hydroxyl group),

(3) a C₂₋₄ alkenyl group,

(4) a C₁₋₄ alkylsulfonyl group or

(5) a C₁₋₄ alkylcarbonyl group

(wherein said C₁₋₄ alkylcarbonyl group is optionally substituted by a hydroxyl group or a C₁₋₄ alkoxy group),

R¹⁴ is a hydrogen atom or a C₁₋₄ alkyl group,

R¹⁸ and R¹⁹ are each independently a hydrogen atom or a C₁₋₄ alkyl group, or may form C₃₋₇ cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom),

A² is a C₁₋₄ alkylene group optionally substituted by a C₃₋₇ cycloalkyl group,

Z is

(1) a C₃₋₇ cycloalkyl group

(wherein said C₃₋₇ cycloalkyl group is optionally substituted by a phenyl group optionally substituted by a halogen atom),

(2) an aryl group

(wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group optionally substituted by a substituent selected from the group consisting of a C₁₋₄ alkyl group and a C₁₋₄ alkylcarbonyl group,

(b) a C₃₋₇ cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,

(c) a carboxy group,

(d) a halogen atom,

(e) a C₁₋₈ alkyl group,

(f) a C₁₋₄ haloalkyl group,

(g) a C₁₋₄ alkylamino group,

(h) a di(C₁₋₄ alkyl) amino group,

(i) a C₁₋₄ alkylthio group and

(j) a C₁₋₄ alkoxy group),

(3) an aromatic heterocyclic group

(wherein said aromatic heterocyclic group is optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group,

(b) a C₁₋₆ alkyl group,

(c) an aryl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group,

(d) a halogen atom,

(e) a C₁₋₄ haloalkyl group,

(f) a carboxy group,

- (g) a C₃₋₇ cycloalkyl group and
- (h) a C₁₋₄ alkoxy group),

- (4) an indanyl group or
- (5) a piperazinyl group

(wherein said piperazinyl group is optionally substituted by a substituent selected from the group consisting of

- (a) a phenyl group,
- (b) a phenyl C₁₋₄ alkyl group,
- (c) a benzoyl group optionally substituted by a halogen atom and
- (d) a phenyl C₁₋₄ alkoxy carbonyl group),

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[3] The azole compound of [2], wherein W is a sulfur atom or an oxygen atom, R is

- (1) -COOR⁷ wherein R⁷ is a hydrogen atom,
 - (2) -X¹-A¹-COOR⁷
- wherein
- X¹ is -O-,
 - A¹ is a C₁₋₄ alkylene group,
 - R⁷ is a hydrogen atom or
 - (3) a tetrazolyl group;

R¹, R², R³ and R⁴ are each independently

- (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a hydroxyl group,
- (4) an optionally substituted C₃₋₇ cycloalkyl C₁₋₄ alkoxy group or
- (5) an optionally substituted aralkoxy group;

A is a group represented by -(CH₂)_m-X-

wherein

X is -N(R⁸)-, -C(R⁹)(R¹⁰)- or -CO-

wherein

R⁸ is a hydrogen atom or a C₁₋₆ alkyl group, wherein said C₁₋₆ alkyl group is optionally substituted by a substituent selected from the group consisting of a C₁₋₄ alkoxy group, an aryloxy group, -N(R¹¹)(R¹²)(R¹¹ and R¹² are each independently a hydrogen atom or a C₁₋₄ alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a C₃₋₇ cycloalkyl group and an optionally substituted aryl group,

R⁹ and R¹⁰ are each independently a hydrogen atom or a C₁₋₄ alkyl group, or may form C₃₋₇ cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, m is 0 or an integer of 1 to 3;

B is an aryl group or an aromatic heterocyclic group;

R⁵ is

- (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a C₁₋₄ alkyl group or
- (4) a C₁₋₄ alkoxy group;

R⁶ is -(Y)_{s1}-(A²)_s-Z

wherein

s₁ and s are each independently 0 or 1,

Y is -O-, -S(O)_t-, -N(R¹³)-, -N(R¹⁴)-CO- or -N(R¹⁴)-SO₂-

wherein

t is 0, 1 or 2,

R¹³ is

- (1) a hydrogen atom,
 (2) a C₁₋₄ alkyl group
 (wherein said C₁₋₄ alkyl group is optionally substituted by a substituent selected from the group consisting of

- (a) a C₃₋₇ cycloalkyl group,
 (b) an optionally substituted aryl group,
 (c) an optionally substituted heterocyclic group and
 (d) a hydroxyl group),

- (3) a C₂₋₄ alkenyl group,
 (4) a C₁₋₄ alkylsulfonyl group or
 (5) a C₁₋₄ alkylcarbonyl group

(wherein said C₁₋₄ alkylcarbonyl group is optionally substituted by a hydroxyl group or a C₁₋₄ alkoxy group),

R¹⁴ is a hydrogen atom or a C₁₋₄ alkyl group,

A² is a C₁₋₄ alkylene group optionally substituted by a C₃₋₇ cycloalkyl group,

Z is

- (1) a C₃₋₇ cycloalkyl group
 (wherein said C₃₋₇ cycloalkyl group is optionally substituted by a phenyl group),
 (2) an aryl group
 (wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group or a C₁₋₄ alkylcarbonyl group,
 (b) a C₃₋₇ cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,
 (c) a carboxy group,
 (d) a halogen atom,
 (e) a C₁₋₈ alkyl group,
 (f) a C₁₋₄ haloalkyl group,
 (g) a C₁₋₄ alkylamino group and
 (h) a di (C₁₋₄ alkyl) amino group,
 (i) a C₁₋₄ alkylthio group and
 (j) a C₁₋₄ alkoxy group),

- (3) an aromatic heterocyclic group
 (wherein said aromatic heterocyclic group is optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group,
 (b) a C₁₋₄ alkyl group and
 (c) a phenyl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group),

- (4) an indanyl group or
 (5) a piperazinyl group

(wherein said piperazinyl group is optionally substituted by a substituent selected from the group consisting of

- (a) a phenyl group,
 (b) a phenyl C₁₋₄ alkyl group and
 (c) a phenyl C₁₋₄ alkoxycarbonyl group),

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[4] The azole compound of [3], wherein W is a sulfur atom and m is 0 or 1, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[5] The azole compound of [4], wherein A is $-(CH_2)_m-X-$ wherein

X is $-N(R^8)-$ wherein R^8 is a hydrogen atom or a C_{1-6} alkyl group, wherein said C_{1-6} alkyl group is optionally substituted by a substituent selected from the group consisting of a C_{1-4} alkoxy group, an aryloxy group, $-N(R^{11})(R^{12})$ (R^{11} and R^{12} are each independently a hydrogen atom or a C_{1-4} alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a C_{3-7} cycloalkyl group and an optionally substituted aryl group, and m is 0 or 1,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[6] The azole compound of [5], wherein R is $-X^1-A^1-COOR^7$ wherein each symbol is as defined in [3], a prodrug thereof or a pharmaceutically acceptable salt thereof.

[7] The azole compound of [5], wherein R is $-COOR^7$ wherein R^7 is a hydrogen atom, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[8] The azole compound of [7], wherein R^1 , R^2 , R^3 and R^4 are hydrogen atoms, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[9] The azole compound of [8], wherein B is a phenyl group, a thiazolyl group, a pyridyl group, a benzothiazolyl group, a benzoimidazolyl group or a benzoxazolyl group,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[10] The azole compound of [9], wherein B is a phenyl group, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[11] The azole compound of [10], wherein R^5 is a hydrogen atom, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[12] The azole compound of [11], wherein, for R^6 , Z is

(1) a C_{3-7} cycloalkyl group

(wherein said C_{3-7} cycloalkyl group is optionally substituted by a phenyl group),

(2) an aryl group

(wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group optionally substituted by a C_{1-4} alkyl group or a C_{1-4} alkylcarbonyl group,

(b) a C_{3-7} cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C_{1-4} alkyl group,

(c) a carboxy group,

(d) a halogen atom,

(e) a C_{1-8} alkyl group,

(f) a C_{1-4} haloalkyl group,

(g) a C_{1-4} alkylamino group,

(h) a di(C_{1-4} alkyl) amino group,

(i) a C_{1-4} alkylthio group and

(j) a C_{1-4} alkoxy group) or

(3) an aromatic heterocyclic group

(wherein said aromatic heterocyclic group is optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group,

(b) a C_{1-4} alkyl group and

(c) a phenyl group optionally substituted by a halogen atom or a C_{1-4} haloalkyl group),

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[13] The azole compound of [12], wherein Z is an aryl group optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group optionally substituted by a C_{1-4} alkyl group or a C_{1-4} alkylcarbonyl group,

(b) a C_{3-7} cycloalkyl group optionally substituted by a substituent selected from the group consisting of a

hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,

(c) a carboxy group,

(d) a halogen atom,

(e) a C₁₋₈ alkyl group,

(f) a C₁₋₄ haloalkyl group,

(g) a C₁₋₄ alkylamino group,

(h) a di(C₁₋₄ alkyl)amino group,

(i) a C₁₋₄ alkylthio group and

(j) a C₁₋₄ alkoxy group,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[14] The azole compound of [13], wherein Z is a phenyl group substituted by a substituent selected from the group consisting of

(a) a cyclohexyl group or a cyclopentyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,

(b) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group or a C₁₋₄ alkylcarbonyl group (wherein said heterocyclic group is selected from the group consisting of a piperidinyl group, a morpholinyl group, a piperazinyl group, a tetrahydropyranyl group, a pyrrolidinyl group and a pyrrolyl group) and

(c) a C₁₋₈ alkyl group,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

[15] The azole compound of [14], wherein Z is a phenyl group substituted by a cyclohexyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[16] The azole compound of [13] or [14] wherein, for R⁶, Y is -O-, -N(R¹³)- or -N(R¹⁴)-CO- wherein

R¹³ is a hydrogen atom, a C₁₋₄ alkyl group or a C₂₋₄ alkenyl group, wherein said C₁₋₄ alkyl group is optionally substituted by a substituent selected from the group consisting of a C₃₋₇ cycloalkyl group, an optionally substituted aryl group, an optionally substituted heterocyclic group and a hydroxyl group, R¹⁴ is a hydrogen atom or a C₁₋₄ alkyl group, and s1 is 1, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[17] The azole compound of [16], wherein, for R⁶, A² is a methylene group, a prodrug thereof or a pharmaceutically acceptable salt thereof.

[18] A pharmaceutical composition comprising an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[19] A pharmaceutical composition for inhibiting protein Tyrosine Phosphatase 1B, which comprises an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[20] A pharmaceutical composition for treating diabetes, which comprises an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[21] A pharmaceutical composition for treating hyperlipidemia, which comprises an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[22] The pharmaceutical composition of [18], which is used in combination with a different therapeutic drug for hyperlipidemia.

[23] The pharmaceutical composition of [22], wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.

[24] The pharmaceutical composition of [23], wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.

[25] The pharmaceutical composition of [18], which is used in combination with a different therapeutic drug for diabetes.

[26] The pharmaceutical composition of [25], which is used in combination with a therapeutic agent for diabetes selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin preparation and an insulin sensitizer.

[27] The pharmaceutical composition of [26], wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glyclopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochloride.

[28] The pharmaceutical composition of [20], which is used in combination with a different therapeutic drug for

hyperlipidemia.

[29] The pharmaceutical composition of [28], wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.

[30] The pharmaceutical composition of [29], wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.

[31] The pharmaceutical composition of [20], which is used in combination with a different therapeutic drug for diabetes.

[32] The pharmaceutical composition of [31], which is used in combination with a therapeutic drug for diabetes selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin preparation and an insulin sensitizer.

[33] The pharmaceutical composition of [32], wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glycopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochloride.

[34] The pharmaceutical composition of [21], which is used in combination with a different therapeutic drug for hyperlipidemia.

[35] The pharmaceutical composition of [34], wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.

[36] The pharmaceutical composition of [35], wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.

[37] The pharmaceutical composition of [21], which is used in combination with a different therapeutic drug for diabetes.

[38] The pharmaceutical composition of [37], which is used in combination with a therapeutic drug for diabetes selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin preparation and an insulin sensitizer.

[39] The pharmaceutical composition of [38], wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glycopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochloride.

[40] A method of inhibiting protein Tyrosine Phosphatase 1B, which comprises administering an effective amount of an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal.

[41] A method of treating diabetes, which comprises administering an effective amount of an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal.

[42] A method of treating hyperlipidemia, which comprises administering an effective amount of an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal.

[43] Use of an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof for the production of a protein tyrosine phosphatase 1B inhibitor.

[44] Use of an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof for the production of a therapeutic agent for diabetes.

[45] Use of an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof for the production of a therapeutic agent for hyperlipidemia.

[46] A commercial package comprising the pharmaceutical composition of [18] and a written matter associated therewith, the written matter stating that said pharmaceutical composition can or should be used for inhibiting protein Tyrosine Phosphatase 1B.

[47] A commercial package comprising the pharmaceutical composition of [18] and a written matter associated therewith, the written matter stating that said pharmaceutical composition can or should be used for treating diabetes.

[48] A commercial package comprising the pharmaceutical composition of [18] and a written matter associated therewith, the written matter stating that said pharmaceutical composition can or should be used for treating hyperlipidemia.

[49] A method of treating hyperlipidemia, which comprises administering an effective amount of an azole compound of any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal, and administering an effective amount of a different therapeutic drug for hyperlipidemia to said mammal.

[50] The method of treating of [49], wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.

[51] The method of treating of [50], wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.

[52] A method of treating diabetes, which comprises administering an effective amount of an azole compound of

any of [1] to [17], a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal, and administering an effective amount of a different therapeutic agent for diabetes to said mammal.

[53] The method of treating of [52], wherein the therapeutic agent for diabetes is selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin preparation and an insulin sensitizer.

[54] The method of treating of [53], wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glycopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochloride.

[0025] The definitions of respective substituents and respective moieties used in the present specification are as follows.

[0026] In the present specification, "C₁₋₆" means that the carbon number is 1 to 6.

[0027] The "halogen atom" is a fluorine atom, a chlorine atom, a bromine atom or an iodine atom, with preference given to a fluorine atom and a chlorine atom.

[0028] The "lower alkyl group" is a linear or branched chain alkyl group having 1 to 6 carbon atoms, which is exemplified by methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, sec-butyl group, tert-butyl group, pentyl group, isopentyl group, neopentyl group, tert-pentyl group, 1-ethylpropyl group and hexyl group. Preferred is a linear or branched chain alkyl group having 1 to 4 carbon atoms.

[0029] Preferably, it is a C₁₋₄ alkyl group for R¹, R², R³, R⁴, R⁵, R⁷, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁸ and R¹⁹, and a C₁₋₆ alkyl group for R⁶, R¹⁶ and R¹⁷.

[0030] The "lower haloalkyl group" is a haloalkyl group wherein a linear or branched chain alkyl group having 1 to 6 carbon atoms is substituted by the above-defined "halogen atom", which is exemplified by fluoromethyl group, difluoromethyl group, trifluoromethyl group, bromomethyl group, chloromethyl group, 1,2-dichloromethyl group, 2,2-dichloromethyl group, 2,2,2-trifluoroethyl group and the like. Preferred is a linear or branched chain haloalkyl group having 1 to 4 carbon atoms, and particularly preferred is trifluoromethyl group.

[0031] Preferably, it is a C₁₋₄ haloalkyl group for R¹, R², R³, R⁴ and R⁵.

[0032] The "lower alkylene group" is a linear or branched chain alkylene group having 1 to 6 carbon atoms, which is exemplified by methylene group, ethylene group, trimethylene group, propylene group, tetramethylene group, pentamethylene group, hexamethylene group and the like. Preferred is a linear or branched chain alkylene group having 1 to 4 carbon atoms and particularly preferred is methylene group.

[0033] Preferably, it is a C₁₋₄ alkylene group for A¹ and A².

[0034] The "lower alkoxy group" is a linear or branched chain alkoxy group having 1 to 6 carbon atoms, which is exemplified by methoxy group, ethoxy group, propoxy group, isopropoxy group, butoxy group, isobutoxy group, tert-butoxy group, pentyloxy group, hexyloxy group and the like. Preferred is a linear or branched chain alkoxy group having 1 to 4 carbon atoms.

[0035] Preferably, it is a C₁₋₄ alkoxy group for R¹, R², R³, R⁴ and R⁵.

[0036] The "lower haloalkoxy group" is a haloalkoxy group wherein a linear or branched chain alkoxy group having 1 to 6 carbon atoms is substituted by the above-defined "halogen atom", which is exemplified by fluoromethyloxy group, difluoromethyloxy group, trifluoromethyloxy group, bromomethyloxy group, chloromethyloxy group, 1,2-dichloromethyloxy group, 2,2-dichloromethyloxy group, 2,2,2-trifluoroethyloxy group and the like. Preferred is a linear or branched chain haloalkoxy group having 1 to 4 carbon atoms, and particularly preferred is a trifluoromethyloxy group.

[0037] Preferably, it is a C₁₋₄ haloalkoxy group for R¹, R², R³ and R⁴.

[0038] The "aryl group" is an aromatic hydrocarbon group having 6 to 14 carbon atoms, which is exemplified by phenyl group, naphthyl group, biphenyl group (e.g., 2-biphenyl group, 3-biphenyl group, 4-biphenyl group etc.), anthryl group and the like. Preferred are a phenyl group and a biphenyl group, and more preferred is a phenyl group.

[0039] Preferably, it is a C₆₋₁₄ aryl group for R¹⁶, R¹⁷, B and Z.

[0040] The "aryloxy group" is an aryloxy group wherein the "aryl moiety" is the above-defined "aryl group", which is exemplified by phenoxy group, naphthyloxy group, biphenyloxy group (e.g., 2-biphenyloxy group, 3-biphenyloxy group, 4-biphenyloxy group), anthryloxy group and the like. Preferred are a phenoxy group and a biphenyloxy group, and more preferred is a phenoxy group.

[0041] The "aralkyloxy group" is an aralkyloxy group wherein the "aryl moiety" is the above-defined "aryl group" and the "alkyl moiety" is a linear or branched chain alkyl group having 1 to 4 carbon atoms, which is exemplified by benzyloxy group, phenethyloxy group, 3-phenylpropyloxy group and the like. Preferred is a benzyloxy group.

[0042] Preferably, it is a C₆₋₁₄ aryl C₁₋₄ alkyloxy group for R¹, R², R³ and R⁴.

[0043] The "lower cycloalkyl group" is a cycloalkyl group having 3 to 7 carbon atoms, which is exemplified by cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group and cycloheptyl group. Preferred is a cycloalkyl group having 5 to 7 carbon atoms, and particularly preferred is a cyclohexyl group.

[0044] Preferably, it is a C₃₋₇ cycloalkyl group for Z.

[0045] The "lower cycloalkylalkyloxy group" is a cycloalkylalkyloxy group wherein the "cycloalkyl moiety" is the above-defined "lower cycloalkyl group" and the "alkyl moiety" is the above-defined "lower alkyl group", which is exemplified by cyclopropylmethyloxy group, cyclobutylmethyloxy group, cyclopentylmethyloxy group, cyclohexylmethyloxy group, cycloheptylmethyloxy group, 2-cyclopropylethyloxy group, 2-cyclobutylethyloxy group, 2-cyclopentylethyloxy group, 2-cyclohexylethyloxy group, 2-cycloheptylethyloxy group, 3-cyclohexylpropyloxy, 4-cyclohexylbutyloxy group and the like. Preferred is a C₃₋₇ cycloalkyl C₁₋₄ alkyloxy group, more preferred is a C₅₋₇ cycloalkyl C₁₋₄ alkyloxy group, and particularly preferred is a cyclohexyl C₁₋₄ alkyloxy group.

[0046] Preferably, it is a C₅₋₇ cycloalkyl C₁₋₄ alkyloxy group for R¹, R², R³ and R⁴.

[0047] The "lower alkenyl group" is a linear or branched chain alkenyl group having 2 to 6 carbon atoms, which is exemplified by vinyl group, 1-propenyl group, allyl group, 1-methyl-2-propenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group, 1-pentenyl group, 2-pentenyl group, 1-hexenyl group, 2-hexenyl group and the like. Preferred is a linear or branched chain alkenyl group having 2 to 4 carbon atoms.

[0048] Preferably, it is a C₂₋₄ alkenyl group for R¹³.

[0049] The "lower alkylsulfonyl group" is an alkylsulfonyl group wherein the "alkyl moiety" is the above-defined "lower alkyl group", which is exemplified by methylsulfonyl group, ethylsulfonyl group, propylsulfonyl group, isopropylsulfonyl group, butylsulfonyl group, isobutylsulfonyl group, sec-butylsulfonyl group, tert-butylsulfonyl group, pentylsulfonyl group, isopentylsulfonyl group, tert-pentylsulfonyl group, hexylsulfonyl group and the like. Preferred is a C₁₋₄ alkylsulfonyl group.

[0050] Preferably, it is a C₁₋₄ alkylsulfonyl group for R¹³.

[0051] The "lower alkylcarbonyl group" is an alkylcarbonyl group wherein the "alkyl moiety" is the above-defined "lower alkyl group", which is exemplified by acetyl group, propionyl group, butyryl group, isobutyl group, valeryl group, isovaleryl group, pivaloyl group, pentanoyl group, hexanoyl group and the like. Preferred is a C₁₋₄ alkylcarbonyl group wherein the "alkyl moiety" is a linear or branched chain alkyl group having 1 to 4 carbon atoms.

[0052] Preferably, it is a C₁₋₄ alkylcarbonyl group for R¹³.

[0053] When R is a group represented by -COOR⁷ or -X¹-A¹-COOR⁷, and R⁷ is a hydrogen atom, this carboxy group may form a salt. As the salt, alkali metal salts (e.g., potassium salt, sodium salt etc.), alkaline earth metal salts (e.g., calcium salt, magnesium salt etc.) and the like can be mentioned. Preferred is an alkali metal salt.

[0054] The tetrazole ring in the tetrazolyl group for R may form an alkali metal salt. As such alkali metal salt, potassium salt, sodium salt and the like can be mentioned.

[0055] The "lower cycloalkane" that may be formed by R⁹ and R¹⁰ together with a carbon atom bonded thereto is cycloalkane having 3 to 7 carbon atoms, which is exemplified by cyclopropane, cyclobutane, cyclopentane, cyclohexane and cycloheptane. Preferred is cycloalkane having 5 to 7 carbon atoms, and particularly preferred is cyclopentane or cyclohexane.

[0056] The "5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom" that may be formed by R⁹ and R¹⁰ together with a carbon atom bonded thereto is preferably a "saturated 5- to 7-membered hetero ring optionally further having 1 to 3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", which is exemplified by tetrahydropyran, thiane and the like, particularly preferably tetrahydropyran.

[0057] The "aromatic heterocyclic group" for B is a "monocyclic or fused 5- to 14-membered aromatic heterocyclic group containing 1-3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", which is exemplified by furyl group, thienyl group, pyrrolyl group, oxazolyl group, isooxazolyl group, thiazolyl group, isothiazolyl group, imidazolyl group, pyrazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, indolyl group, isoindolyl group, benzofuranyl group, benzothienyl group, benzoimidazolyl group, benzothiazolyl group, benzoxazolyl group, indolizynyl group, quinolyl group, isoquinolyl group, quinazolinyl group, cinnolinyl group, quinoxalinyl group, phthalazinyl group, acridinyl group, phenazinyl group, naphthyridinyl group and the like. Preferred is a "monocyclic or fused 5- to 10-membered aromatic heterocyclic group containing 1 to 3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", and furyl group, thienyl group, pyrrolyl group, oxazolyl group, isooxazolyl group, thiazolyl group, isothiazolyl group, imidazolyl group, pyrazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, indolyl group, isoindolyl group, benzofuranyl group, benzothienyl group, benzoimidazolyl group, benzothiazolyl group, benzoxazolyl group and the like can be mentioned. Particularly preferred are thiazolyl group, pyridyl group, benzothiazolyl group, benzoimidazolyl group and benzoxazolyl group.

[0058] The "lower cycloalkylalkyloxy group" for R¹, R², R³ and R⁴ is optionally substituted by 1 to 3 substituents selected from the following. As such substituent, halogen atom, C₁₋₄ alkyl group, C₁₋₄ haloalkyl group, C₁₋₄ alkoxy group, carboxy group, hydroxyl group, cyano group, nitro group, amino group, alkoxycarbonyl group (alkoxy moiety has 1 to 4 carbon atoms) and the like can be mentioned.

[0059] The "optionally substituted lower cycloalkylalkyloxy group" for R¹, R², R³ and R⁴ is preferably 2-cyclohexylethyloxy group.


[0060] The "aralkyloxy group" for R^1 , R^2 , R^3 and R^4 is optionally substituted by 1 to 3 substituents selected from the following. As such substituent, halogen atom, C_{1-4} alkyl group, C_{1-4} haloalkyl group, C_{1-4} alkoxy group, carboxy group, hydroxyl group, cyano group, nitro group, amino group, alkoxycarbonyl (alkoxy moiety has 1 to 4 carbon atoms) group and the like can be mentioned. Preferable substituent is carboxy group.

[0061] The "optionally substituted aralkyloxy group" for R^1 , R^2 , R^3 and R^4 is preferably benzyloxy group, carboxy-benzyloxy group and the like.


[0062] The "lower alkyl group" for R^8 is optionally substituted by a substituent selected from the group consisting of a lower alkoxy group, an aryloxy group, $-N(R^{11})(R^{12})$ (R^{11} and R^{12} are each independently a hydrogen atom or a lower alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a lower cycloalkyl group and an optionally substituted aryl group.

[0063] The "optionally substituted aryl group" which is a substituent on the "lower alkyl group" for R^8 is optionally substituted by 1 to 3 substituents selected from the following. As such substituent, halogen atom, C_{1-4} alkyl group, C_{1-4} haloalkyl group, C_{1-4} alkoxy group, carboxy group, hydroxyl group, cyano group, nitro group, amino group, alkoxycarbonyl group (alkoxy moiety has 1 to 4 carbon atoms) and the like can be mentioned. Preferable substituents are halogen atom and C_{1-4} haloalkyl group.

[0064] The "5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom" that may be formed by R^{11} and R^{12} together with the nitrogen atom bonded thereto is preferably "a saturated or unsaturated 5- to 7-membered hetero ring optionally further having 1-3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", which is exemplified by a hetero ring selected from the group consisting of



particularly preferably



[0065] The "lower alkyl group" for R^{13} is optionally substituted by a substituent selected from the group consisting of C_{3-7} cycloalkyl group, optionally substituted aryl group, optionally substituted heterocyclic group and hydroxyl group.

[0066] The "optionally substituted aryl group" which is a substituent on the "lower alkyl group" for R^{13} is optionally substituted by 1 to 3 substituents selected from the following. As such substituent, halogen atom, C_{1-4} alkyl group, C_{1-4} haloalkyl group, C_{1-4} alkoxy group, carboxy group, hydroxyl group, cyano group, nitro group, amino group, alkoxycarbonyl group (alkoxy moiety has 1 to 4 carbon atoms) and the like can be mentioned. Preferable substituent is a halogen atom or C_{1-4} haloalkyl group.

[0067] The "optionally substituted heterocyclic group" which is a substituent on the "lower alkyl group" for R^{13} is preferably a "saturated or unsaturated 5- to 7-membered heterocyclic group containing 1 to 3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", which is exemplified by furyl group, thienyl group, pyrrolyl group, oxazolyl group, isooxazolyl group, thiazolyl group, isothiazolyl group, imidazolyl group, pyrazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, tetrahydrofuryl group, tetrahydrothienyl group, pyrrolidinyl group, pyrazolidinyl group, imidazolidinyl group, oxazolidinyl group, thiazolidinyl group, tetrahydropyranyl group, dioxanyl group, piperidinyl group, piperazinyl group, morpholinyl group and the like, with preference given to tetrahydropyranyl group.

[0068] The "optionally substituted hetero ring" which is a substituent on the "lower alkyl group" for R^{13} is optionally substituted by 1 to 3 substituents selected from the following. As such substituent, halogen atom, C_{1-4} alkyl group, C_{1-4} haloalkyl group, C_{1-4} alkoxy group, carboxy group, hydroxyl group, cyano group, nitro group, amino group, alkoxycarbonyl group (alkoxy moiety has 1 to 4 carbon atoms) and the like can be mentioned.

[0069] The "lower alkylcarbonyl group" for R^{13} is optionally substituted by a hydroxyl group or a lower alkoxy group.

[0070] As the lower alkoxy group which is a substituent on the "lower alkylcarbonyl group" for R^{13} , the above-defined "lower alkoxy group" can be mentioned, with preference given to C_{1-4} alkoxy group.


[0071] The "lower cycloalkane" that may be formed by R^{18} and R^{19} together with a carbon atom bonded thereto is cycloalkane having 3 to 7 carbon atoms, which is exemplified by cyclopropane, cyclobutane, cyclopentane, cyclohex-

ane and cycloheptane. It is preferably cycloalkane having 5 to 7 carbon atoms, and particularly preferably cyclopentane or cyclohexane.

[0072] The "5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom" that may be formed by R¹⁸ and R¹⁹ together with a carbon atom bonded thereto is preferably, a "saturated 5- to 7-membered hetero ring optionally further having 1 to 3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", which is exemplified by tetrahydropyran, thiane and the like, particularly preferably tetrahydropyran.

[0073] The lower alkylene group for A² is optionally substituted by a lower cycloalkyl group. As such lower cycloalkyl group, cycloalkyl group having 3 to 7 carbon atoms can be mentioned, which is exemplified by cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group and cycloheptyl group. It is preferably a cycloalkyl group having 5 to 7 carbon atoms, particularly preferably a cyclohexyl group.

[0074] The "lower alkylene group optionally substituted by a lower cycloalkyl group" for A² is preferably a "C₁₋₄ alkylene group optionally substituted by C₃₋₇ cycloalkyl group", more preferably,



[0075] The "lower cycloalkyl group" for Z is preferably a C₃₋₇ cycloalkyl group, more preferably a cyclopentyl group or a cyclohexyl group, still more preferably a cyclohexyl group.


[0076] The "lower cycloalkyl group" for Z may be substituted by (a) a halogen atom, (b) a C₁₋₆ alkyl group, (c) a C₁₋₄ haloalkyl group, (d) a carboxy group, (e) a C₃₋₇ cycloalkyl group, (f) a C₁₋₄ alkoxy group, (g) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group or (h) a phenyl group, wherein said phenyl group is further optionally substituted by 1 to 5 (preferably 1 to 3) halogen atoms. Such substituent of the "lower cycloalkyl group" is preferably a phenyl group optionally substituted by 1 to 3 halogen atoms, more preferably a phenyl group.

[0077] The "aryl group" for Z is preferably a phenyl group or a biphenyl group (e.g., 2-biphenyl group, 3-biphenyl group, 4-biphenyl group), more preferably a phenyl group.


[0078] The "aryl group" for Z is optionally substituted by 1 to 5 (preferably 1 to 3) substituents selected from the following;

- (a) a heterocyclic group optionally substituted by a substituent selected from the group consisting of a lower alkyl group and a lower alkylcarbonyl group,
- (b) a lower cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a lower alkyl group,
- (c) a carboxy group,
- (d) a halogen atom,
- (e) an alkyl group,
- (f) a lower haloalkyl group,
- (g) a lower alkylamino group,
- (h) a di(lower alkyl)amino group,
- (i) a lower alkylthio group and
- (j) a lower alkoxy group.

[0079] The "heterocyclic group" of the "heterocyclic group optionally substituted by a substituent selected from the group consisting of lower alkyl group and lower alkylcarbonyl group" is preferably a "saturated or unsaturated 5- to 7-membered heterocyclic group containing 1 to 3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", which is exemplified by furyl group, thienyl group, pyrrolyl group, oxazolyl group, isoxazolyl group, thiazolyl group, isothiazolyl group, imidazolyl group, pyrazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, tetrahydrofuryl group, tetrahydrothienyl group, pyrrolidinyl group, pyrazolidinyl group, imidazolidinyl group, oxazolidinyl group, thiazolidinyl group, tetrahydropyran group, dioxanyl group, piperidinyl group, piperazinyl group, morpholinyl group and the like. It is preferably piperidinyl group, morpholinyl group, piperazinyl group, pyrrolidinyl group, pyrrolyl group or tetrahydropyran group, more preferably a group selected from the group consisting of



particularly preferably a group selected from the group consisting of



The substituent on said "heterocyclic group" is preferably a C_{1-4} alkyl group or a C_{1-4} alkylcarbonyl group (alkyl moiety has 1 to 4 carbon atoms).

[0080] The "lower cycloalkyl" of the "lower cycloalkyl group optionally substituted by a substituent selected from the group consisting of hydroxyl group, an oxo group, a halogen atom and lower alkyl group" is preferably C_{3-7} cycloalkyl group, more preferably cyclohexyl group. The "lower cycloalkyl group" is optionally substituted by 1 to 5 (preferably 1 to 3) substituents selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a lower alkyl group. The substituent on the "lower cycloalkyl group" is preferably a hydroxyl group, an oxo group, a halogen atom or a C_{1-4} alkyl group.

[0081] As the "halogen atom" which is a substituent on "aryl group" for Z, the above-defined "halogen atom" can be preferably mentioned, such as fluorine atom, chlorine atom and bromine atom.

[0082] The "alkyl group" which is a substituent on the "aryl group" for Z is preferably a linear or branched chain alkyl group having 1 to 8 carbon atoms, which is exemplified by methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, sec-butyl group, tert-butyl group, pentyl group, isopentyl group, neopentyl group, tert-pentyl group, 1-ethylpropyl group, hexyl group, heptyl group, 1-propylbutyl group, octyl group and the like.

[0083] The "lower haloalkyl group" which is a substituent on the "aryl group" for Z is the above-defined "lower haloalkyl group", which is preferably a C_{1-4} haloalkyl group.

[0084] The "lower alkylamino group" which is a substituent on the "aryl group" for Z is an alkylamino group wherein the "alkyl moiety" is the above-defined "lower alkyl group", which is exemplified by methylamino group, ethylamino group, propylamino group, isopropylamino group, butylamino group, isobutylamino group, sec-butylamino group, tert-butylamino group, pentylamino group, isopentylamino group, neopentylamino group, tert-pentylamino group, hexylamino group and the like. It is preferably a C_{1-4} alkylamino group.

[0085] The "di(lower alkyl)amino group" which is a substituent on the "aryl group" for Z is dialkylamino group wherein the "alkyl moiety" is the above-defined "lower alkyl group", which is exemplified by dimethylamino group, diethylamino group, dipropylamino group, diisopropylamino group, dibutylamino group, diisobutylamino group, di(sec-butyl)amino group, di(tert-butyl)amino group, dipentylamino group, diisopentylamino group, di(tert-pentyl)amino group, dihexylamino group, N-ethyl-N-methylamino group, N-methyl-N-propylamino group, N-ethyl-N-propylamino group and the like. Preferred is a di(C_{1-4} alkyl) amino group.

[0086] The "lower alkylthio group" which is a substituent on the "aryl group" for Z is alkylthio group wherein the "alkyl moiety" is the above-defined "lower alkyl group", which is exemplified by methylthio group, ethylthio group, propylthio group, isopropylthio group, butylthio group, isobutylthio group, sec-butylthio group, tert-butylthio group, pentylthio group, isopentylthio group, neopentylthio group, tert-pentylthio group, hexylthio group and the like. It is preferably a C_{1-4} alkylthio group.

[0087] The "lower alkoxy group" which is a substituent on the "aryl group" for Z is the above-defined "lower alkoxy group", preferably a C_{1-4} alkoxy group.

[0088] The "aromatic heterocyclic group" for Z is preferably a "monocycle or fused 5- to 10-membered aromatic heterocyclic group containing 1-3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", and furyl group, thienyl group, pyrrolyl group, oxazolyl group, isooxazolyl group, thiazolyl group, isothiazolyl group, imidazolyl group, pyrazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, indolyl group, isoindolyl group, benzofuranyl group, benzothienyl group, benzoimidazolyl group, benzothiazolyl group, benzoxazolyl group and the like can be mentioned. It is particularly preferably a thiazolyl group or a pyridyl group.


[0089] The "aromatic heterocyclic group" for Z is optionally substituted by 1 to 5 (preferably 1 to 3) substituents selected from the following;

(a) a heterocyclic group optionally substituted by C_{1-4} alkyl group,


- (b) a C₁₋₆ alkyl group,
- (c) an aryl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group,
- (d) a halogen atom,
- (e) a C₁₋₄ haloalkyl group,
- (f) a carboxy group,
- (g) a C₃₋₇ cycloalkyl group and
- (h) a C₁₋₄ alkoxy group.

[0090] Such substituent is preferably (a) a heterocyclic group, (b) a C₁₋₆ alkyl group or (c) an aryl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group.

[0091] The "heterocyclic group" which is a substituent on the "aromatic heterocyclic group" for Z is preferably a "saturated or unsaturated 5- to 7-membered heterocyclic group containing 1 to 3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom", which is exemplified by furyl group, thienyl group, pyrrolyl group, oxazolyl group, isooxazolyl group, thiazolyl group, isothiazolyl group, imidazolyl group, pyrazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, tetrahydrofuryl group, tetrahydrothienyl group, pyrrolidinyl group, pyrazolidinyl group, imidazolidinyl group, oxazolidinyl group, thiazolidinyl group, tetrahydropyranyl group, dioxan-yl group, piperidinyl group, piperazinyl group, morpholinyl group and the like. It is preferably piperidinyl group, morpholinyl group, piperazinyl group, tetrahydropyranyl group, pyrrolidinyl group or pyrrolyl group, more preferably a group selected from the group consisting of



particularly preferably



[0092] The "aryl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group" which is a substituent on the "aromatic heterocyclic group" for Z is preferably a "phenyl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group".

[0093] The "piperazinyl group" for Z is optionally substituted by 1 to 5 (preferably 1 to 3) substituents selected from the following:

- (a) a phenyl group,
- (b) a phenyl lower alkyl group,
- (c) a benzoyl group optionally substituted by a halogen atom and
- (d) a phenyl lower alkoxy carbonyl group.

[0094] The "phenyl lower alkyl group" which is a substituent on the "piperazinyl group" for Z is a phenylalkyl group wherein the "alkyl moiety" is the above-defined "lower alkyl group", which is exemplified by benzyl group, phenethyl group, 1-phenylethyl group, 3-phenylpropyl group and the like. Preferably, it is a phenyl C₁₋₄ alkyl group.

[0095] The "benzoyl group optionally substituted by a halogen atom" which is a substituent on the "piperazinyl group" for Z is preferably a benzoyl group optionally substituted by 1 to 5 the above-defined "halogen atoms", which is exemplified by chlorobenzoyl group, bromobenzoyl group and the like.

[0096] The "phenyl lower alkoxy carbonyl group" which is a substituent on the "piperazinyl group" for Z is a phenylalkoxy carbonyl group wherein the "alkoxy moiety" is the above-defined "lower alkoxy group", which is exemplified by a benzyloxy carbonyl group and the like. Preferably, it is a phenyl C₁₋₄ alkoxy carbonyl group.

[0097] In the formula [I], preferable substituents are as follows.

W is preferably a sulfur atom.

R is preferably -COOR⁷ wherein R⁷ is a hydrogen atom.

R¹, R², R³ and R⁴ are preferably hydrogen atoms.

A is preferably -(CH₂)_m-X-

wherein

X is a group represented by $-N(R^8)-$ wherein R^8 is a hydrogen atom or a C_{1-6} alkyl group, wherein said C_{1-6} alkyl group is optionally substituted by a substituent selected from the group consisting of a C_{1-4} alkoxy group, an aryloxy group, $-N(R^{11})(R^{12})$ (R^{11} and R^{12} are each independently a hydrogen atom or a C_{1-4} alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a C_{3-7} cycloalkyl group and an optionally substituted aryl group, and m is 0 or an integer of 1 to 3.

B is preferably a phenyl group, a thiazolyl group, a pyridyl group, a benzothiazolyl group, a benzoimidazolyl group or a benzoxazolyl group, more preferably a phenyl group.

R^5 is preferably a hydrogen atom.

Z is preferably

(1) a C_{3-7} cycloalkyl group

(wherein said C_{3-7} cycloalkyl group is optionally substituted by a phenyl group optionally substituted by a halogen atom),

(2) an aryl group

(wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group optionally substituted by a C_{1-4} alkyl group or a C_{1-4} alkylcarbonyl group,

(b) a C_{3-7} cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C_{1-4} alkyl group,

(c) a carboxy group,

(d) a halogen atom,

(e) a C_{1-8} alkyl group,

(f) a C_{1-4} haloalkyl group,

(g) a C_{1-4} alkylamino group,

(h) a di(C_{1-4} alkyl)amino group,

(i) a C_{1-4} alkylthio group and

(j) a C_{1-4} alkoxy group) or

(3) an aromatic heterocyclic group

(wherein said aromatic heterocyclic group is optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group,

(b) a C_{1-4} alkyl group and

(c) a phenyl group optionally substituted by a halogen atom or a C_{1-4} haloalkyl group).

[0098] Z is more preferably an aryl group optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group optionally substituted by a C_{1-4} alkyl group or a C_{1-4} alkylcarbonyl group,

(b) a C_{3-7} cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C_{1-4} alkyl group,

(c) a carboxy group,

(d) a halogen atom,

(e) a C_{1-8} alkyl group,

(f) a C_{1-4} haloalkyl group,

(g) a C_{1-4} alkylamino group,

(h) a di(C_{1-4} alkyl) amino group and

(i) a C_{1-4} alkylthio group.

[0099] Z is further preferably a phenyl group substituted by a substituent selected from the group consisting of

(a) a cyclohexyl group or a cyclopentyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C_{1-4} alkyl group,

(b) a heterocyclic group optionally substituted by a C_{1-4} alkyl group or a C_{1-4} alkylcarbonyl group (wherein said heterocyclic group is selected from the group consisting of a piperidinyl group, a morpholinyl group, a piperazinyl group, a tetrahydropyranyl group, a pyrrolidinyl group and a pyrrolyl group) and

(c) C₁₋₈ alkyl groups.

[0100] Z is particularly preferably a phenyl group substituted by a cyclohexyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group.

[0101] For R⁶, Y is preferably -O-, -N(R¹³)- or -N(R¹⁴)-CO- wherein

R¹³ is a hydrogen atom, a C₁₋₄ alkyl group or a C₂₋₄ alkenyl group,

wherein said C₁₋₄ alkyl group is optionally substituted by a substituent selected from the group consisting of a C₃₋₇ cycloalkyl group, an optionally substituted aryl group and an optionally substituted heterocyclic group,

R¹⁴ is a hydrogen atom or a C₁₋₄ alkyl group, and s1 is 0 or 1.

[0102] A² is preferably a methylene group.





[0103] The "pharmaceutically acceptable salt" may be any salt as long as it forms a non-toxic salt with the compound represented by the above-mentioned formula [I], and can be obtained by reacting with inorganic acids such as hydrochloric acid, sulfuric acid, phosphoric acid, hydrobromic acid and the like; organic acids such as oxalic acid, malonic acid, citric acid, fumaric acid, lactic acid, malic acid, succinic acid, tartaric acid, acetic acid, trifluoroacetic acid, gluconic acid, ascorbic acid, methylsulfonic acid, benzenesulfonic acid and the like; inorganic bases such as sodium hydroxide, potassium hydroxide, calcium hydroxide, magnesium hydroxide, ammonium hydroxide and the like; organic bases such as methylamine, diethylamine, triethylamine, triethanolamine, ethylenediamine, tris(hydroxymethyl)methylamine, guanidine, choline, cinchonine, N-methyl-D-glucamine and the like; or amino acids such as lysin, histidine, arginine, alanine and the like. The present invention also encompasses water-containing product, hydrate and solvate of each compound.



[0104] The compound represented by the above-mentioned formula [I] contains various isomers. For example, E-form and Z-form are present as geometric isomers, and when an asymmetric carbon atom is present, enantiomer and diastereomer are present as stereoisomers based thereon. In some cases, a tautomer may be present. Accordingly, the present invention encompasses all of these isomers and mixtures thereof.

[0105] The present invention encompasses prodrug and metabolite of the compound of the formula. [I].

[0106] A "prodrug" is a derivative of the compound of the present invention, which has a chemically or metabolically decomposable group, and, after administration to a living organism, restores to the original compound to show the inherent efficacy. It includes a complex free of covalent bond and salt.

[0107] For example, known ester derivatives can be used as prodrugs in the field of pharmaceutical drugs. To be specific, ester derivatives wherein R is a group represented by the following formula can be mentioned.





[0108] When the compound of the present invention is used as a pharmaceutical preparation, it is generally admixed with pharmaceutically acceptable carrier, excipient, diluent, extender, disintegrant, stabilizer, preservative, buffer, emulsifier, aromatic, coloring agent, sweetening agent, thickening agent, corrigent, dissolution aids and other additives, which are known per se and specifically exemplified by water, vegetable oil, alcohols such as ethanol, benzyl alcohol etc., polyethylene glycol, glycerol triacetate, gelatin, lactose, carbohydrate such as starch etc., magnesium stearate, talc, lanolin, vaseline and the like, and prepared into the form of tablet, pill, powder, granule, suppository, injection, eye drop, liquid, capsule, troche, aerosol, elixir, suspension, emulsion, syrup and the like by a conventional method, which can be administered systemically or topically, and orally or parenterally.

[0109] The dose of the compound of the present invention varies depending on the age, body weight, symptom, disease to be treated, administration method and the like, but it is generally administered in the range of 50 mg to 800 mg per dose for an adult once to several times a day.

[0110] The compound [I] of the present invention can be administered to mammals (human, mouse, rat, rabbit, dog, cat, bovine, pig, monkey etc.) as a PTP1B inhibitor, a drug for the prophylaxis or treatment of diabetes, a drug for the prophylaxis or treatment of diabetic complications (retinopathy, nephropathy, neuropathy, cardiac infarction and cerebral infarction based on arteriosclerosis, etc.), a drug for the prophylaxis or treatment of hyperlipidemia, a drug for the prophylaxis or treatment of obesity, neurodegenerative disease and the like or a drug for the prophylaxis or treatment of a disease mediated by PTP1B.

[0111] The compound [I] of the present invention can be administered to a mammal together with other therapeutic agents for diabetes for the purpose of prophylaxis or treatment of diabetes or diabetic complications. In the present invention, "therapeutic agents for diabetes" include therapeutic agents for diabetic complications. In addition, the compound [I] of the present invention can be administered to a mammal together with other therapeutic agents for hyperlipidemia for the purpose of prophylaxis or treatment of hyperlipidemia.

[0112] In the case of combined administration, the compound of the present invention can be administered simultaneously with a different therapeutic agent for diabetes or a different therapeutic drug for hyperlipidemia (hereinafter combination drug) or administered at certain time intervals. In the case of combined administration, a pharmaceutical composition containing the compound of the present invention and a combination drug can be administered. Alternatively, a pharmaceutical composition containing the compound of the present invention and a pharmaceutical composition containing a combination drug may be administered separately. The administration route of each pharmaceutical composition may be the same or different.

[0113] In the case of a combined administration, the compound of the present invention can be administered once to several times a day in the range of 50 mg to 800 mg per dose, or may be administered at a lower dose. A combination drug can be administered at a dose generally employed for the prophylaxis or treatment of diabetes or for the prophylaxis or treatment of hyperlipidemia, or may be administered at a lower dose.

[0114] As other therapeutic agents for diabetes to be used for the combined administration, insulin secretagogue, sulfonylurea, sulfonamide, biguanide, α glucosidase inhibitor, insulin preparation, insulin sensitizer and the like can be mentioned. For example, nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glyclopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin, pioglitazone hydrochloride and the like can be used for combined administration with the compound of the present invention.

[0115] As other therapeutic drugs for hyperlipidemia to be used for the combined administration, statin drugs can be mentioned. For example, lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, cerivastatin and the like can be used for combined administration with the compound of the present invention.

[0116] Now, one example of the production methods of the compounds used to practice the present invention is shown below. However, the production method of the compound of the present invention is not limited to the example.

[0117] Even in the absence of description in the production methods, a protecting group may be introduced into functional groups as necessary and deprotected at a suitable stage, the order of each production method and step is exchanged and the like to efficiently perform production.


[0118] The treatment after reaction in each step may be performed according to a method generally employed, and conventional methods such as isolation and purification, crystallization, recrystallization, silica gel chromatography, preparative HPLC and the like are appropriately selected, or used in combination.

Production Method 1

[0119] In this Production Method, compound [I] wherein W is a sulfur atom is produced.

Production Method 1

[0120]



wherein R_A is $-\text{COOR}^{7'}$ or $-\text{X}^1-\text{A}^1-\text{COOR}^{7'}$ ($R^{7'}$ is a lower alkyl group), Hal is a halogen atom such as a bromine atom, a chlorine atom and the like, and other symbols are as defined above.

Step 1

[0121] The compound [2] can be obtained by reacting compound [1] with a thiocarbonylation agent such as a Lawesson's reagent, phosphorus pentasulfide and the like in a solvent. As the solvent, tetrahydrofuran (THF), 1,2-dimethoxyethane (DME), toluene, xylene, chloroform, dichloromethane, dioxane and the like, or a mixed solvent thereof can be mentioned. The reaction temperature is preferably 50°C - 100°C .

Step 2

[0122] The compound [I]-1 can be obtained by reacting compound [2] with compound [3] in the presence or absence of a base in a solvent under heating. As the solvent, acetonitrile, alcohols (methanol, ethanol, isopropyl alcohol etc.), THF, DME, dioxane and the like, or a mixed solvent thereof can be mentioned. As the base, sodium hydrogen carbonate, potassium hydrogen carbonate and the like can be mentioned.

Production Method 2

[0123] In this Production Method, compound [I] wherein W is an oxygen atom is produced.


5 Production Method 2

[0124]

10

15

20



wherein each symbol is as defined above.

25

Step 3

[0125] The compound [I]-2 can be obtained by reacting compound [1] with compound [3] in a solvent under heating. As the solvent, acetonitrile, alcohols (methanol, ethanol, isopropyl alcohol etc.), xylene, toluene and the like, or a mixed solvent thereof can be mentioned.

30

Production Method 3

[0126] In this Production Method, compound [I] wherein W is a sulfur atom and A is $-(CH_2)_m-N(R^8)-$ is produced.

35

40


45

50

55

Production Method 3

[0127]



wherein $\text{R}^{8'}$ is $-\text{SO}_2\text{R}^{16}$ or a lower alkyl group, wherein said lower alkyl group is optionally substituted by a substituent selected from the group consisting of a lower alkoxy group, an aryloxy group, $-\text{N}(\text{R}^{11})(\text{R}^{12})$, a carboxy group, a lower cycloalkyl group and an optionally substituted aryl group, L is a leaving group such as an iodine atom, a bromine atom, a chlorine atom and the like, and other symbols are as defined above.

Step 4

[0128] The compound [6] can be obtained by reacting compound [4] with 1,1'-thiocarbonyldiimidazole, thiophosgene and the like in a solvent in the presence or absence of a base, then reacting with ammonia. As the solvent, chloroform, dichloromethane, dichloroethane, THF, DME, dioxane, toluene and the like, or a mixed solvent thereof can be mentioned. As the base, triethylamine, diisopropylethylamine, 1,8-diazabicyclo[5.4.0]undeca-7-ene (DBU), sodium hydride and the like can be mentioned. The reaction temperature is preferably -20°C - 50°C .

Step 5

[0129] The compound [I]-3 can be obtained by reacting compound [6] with compound [3] in a solvent under heating in the presence or absence of a base. As the solvent, acetonitrile, alcohols (methanol, ethanol, isopropyl alcohol etc.), THF, DME, dioxane and the like, or a mixed solvent thereof can be mentioned. As the base, sodium hydrogen carbonate, potassium hydrogen carbonate and the like can be mentioned.

Step 6


[0130] The compound [I]-4 can be obtained by reacting compound [I]-3 with compound [7] in a solvent in the presence of a base. As the solvent, dimethylformamide, dimethylacetamide, THF, DME, dioxane, hexamethylphosphoramide (HMPA), dimethyl sulfoxide (DMSO) and the like, or a mixed solvent thereof can be mentioned. As the base, sodium hydride, potassium carbonate, sodium carbonate and the like can be mentioned. The reaction temperature is preferably 0°C - 100°C.

Production Method 4

[0131] In this Production Method, compound [I] wherein R⁷ is a hydrogen atom when R is -COOR⁷ or -X¹-A¹-COOR⁷, or a salt thereof is produced.

Production Method 4

[0132]



wherein R_B is -COOH or -X¹-A¹-COOH, R_C is -COOM or -X¹-A¹-COOM (M is an alkali metal), and other symbols are as defined above.

Step 7

[0133] The compound [I]-6 can be obtained by hydrolyzing compound [I]-5. Hydrolysis can be performed according to conventional methods. For example, a method comprises hydrolyzing in a solvent in the presence of an acid (including a Lewis acid) or a base can be mentioned. As the solvent, alcohols (methanol, ethanol, isopropyl alcohol etc.), tetrahydrofuran, dioxane, DME, N,N-dimethylformamide (DMF), DMSO, water and the like, or a mixed solvent thereof can be mentioned. As the acid, hydrochloric acid, trifluoroacetic acid, sulfuric acid and the like can be mentioned. As the base, alkali metal hydroxide (sodium hydroxide, potassium hydroxide etc.), potassium carbonate, sodium carbonate and the like can be mentioned. The reaction temperature is not particularly limited, and the reaction can be carried out under cooling to under heating.

Step 8


[0134] The compound [I]-7 can be obtained by reacting compound [I]-6 with alkali metal hydroxide [8], according to conventional methods. Alkali metal hydroxide includes sodium hydroxide, potassium hydroxide and the like. This Step can be conducted in a solvent. As the solvent, alcohols (methanol, ethanol etc.), tetrahydrofuran, dioxane, DME and the like, or a mixed solvent thereof can be mentioned. The reaction temperature is not particularly limited, and the reaction can be carried out under cooling to under heating.

Production Method 5

[0135] In this Production Method, compound [I] wherein R is a tetrazolyl group is produced.

5 **Production Method 5**

[0136]



wherein each symbol is as defined above.

20

Step 9

[0137] The compound [I]-8 can be obtained by reacting compound [9] produced in the same manner as in the aforementioned Production Method 1-3 with trimethylsilylazide and dibutyltin oxide in a solvent. As the solvent, toluene, xylene, benzene and the like, or a mixed solvent thereof can be mentioned. The reaction temperature is preferably 50°C - 150°C.

25

Production Method 6

[0138] In this Production Method, compound [I] wherein R⁶ is -N(R¹³)-CH₂-Z is produced.

35

40


45

50

55

Production Method 6

[0139]



wherein each symbol is as defined above.

Step 10

[0140] The compound [11] can be obtained by reducing compound [10]. Reduction can be performed according to conventional methods. For example, compound [10] is subjected to catalytic reduction in a solvent in the presence of a catalyst under a hydrogen atmosphere. As the solvent, alcohols (methanol, ethanol, isopropyl alcohol etc.), tetrahydrofuran, acetic acid and the like can be mentioned. As the catalyst, palladium catalysts such as palladium-carbon etc. and the like can be mentioned.

Step 11

[0141] The compound [I]-9 can be obtained by reacting compound [11] with compound [12] in the presence of a reducing agent. As the reducing agent, sodium triacetoxyborohydride, sodium cyanoborohydride (NaBH_3CN) and the like can be mentioned. The reaction temperature is preferably 0°C - 40°C .

Step 12

[0142] The compound [I]-10 can be obtained by reacting compound [I]-9 with compound [13] in a solvent in the presence of a base. This reaction can be carried out in the same manner as in Step 6 of Production Method 3.

[0143] The Production Methods described in the present specification are examples of the production methods of the compounds of the present invention, and compounds other than those explained in the above can be also produced by combining conventional methods known in the field of organic synthetic chemistry.

Examples


[0144] The compound represented by the formula [I] and Production Methods of the present invention are explained in detail by referring to Production Examples and Examples, which are not to be construed as limitative.

Production Example 1

4-cyclohexylbenzaldehyde

(1) 4-cyclohexylbenzyl alcohol

[0145]




[0146] Under a nitrogen stream, tetrahydrofuran (1.2 L, 15.0 v/w) was added to 4-cyclohexylbenzoic acid (80.0 g, 0.382 mol) and then isobutyl chlorocarbonate (52.0 ml, 0.401 mol) was added. Triethylamine (56.0 ml, 0.401 mol) was added to the reaction mixture with stirring under ice-cooling and the mixture was stirred at the same temperature for 30 min. The resulting precipitate was collected by filtration. To a suspension of sodium borohydride (58.0 g, 1.53 mol) in tetrahydrofuran (160 ml, 2.0 v/w) prepared in a different reaction vessel was carefully added the above-mentioned filtrate with stirring under ice-cooling under a nitrogen stream. After stirring at room temperature for 1.5 hr, distilled water (160 ml, 2.0 v/w) was added with stirring under ice-cooling. After stirring under ice-cooling for 20 min, 2N-hydrochloric acid (825 ml, 4.3 eq) was added. After stirring at room temperature for 30 min, the mixture was extracted with ethyl acetate (400 ml) and the organic layer was washed successively with distilled water (100 ml), 2N-aqueous sodium hydroxide solution (100 ml), distilled water (100 ml) and saturated brine (100 ml), and dried over magnesium sulfate (70 g). After filtration and solvent evaporation, the residue was dried in vacuo to give the title compound (63.8 g, yield 87.8%) as a white solid.

¹H-NMR(300MHz,DMSO-d₆) δ 1.23-1.41(5H,m), 1.67-1.78(5H,m), 2.47(1H,m), 4.43(2H,s), 5.04(1H,brs), 7.15(2H,d, J=8.0Hz), 7.21(2H,d,J=8.0Hz).

(2) 4-cyclohexylbenzaldehyde

[0147]



[0148] To a solution of 4-cyclohexylbenzyl alcohol (121.5 g, 0.639 mol) obtained in Production Example 1(1) in dimethyl sulfoxide (500 ml) was added triethylamine (249 ml, 1.79 mol). Pyridine-sulfur trioxide complex (163 g, 1.02 mol) was gradually added with stirring under ice-cooling, and the mixture was stirred at room temperature for 2 hr. Water (500 ml) was added dropwise to the reaction mixture with stirring under ice-cooling, and the mixture was extracted with a mixed solvent (1:1) of n-hexane and ethyl acetate, washed with saturated brine and dried over sodium sulfate. After filtration and solvent evaporation, the residue was dried in vacuo to give the title compound (112 g, yield 93.4%) as a colorless oil.


¹H-NMR(300MHz,CDCl₃) δ 1.20-1.53(5H,m), 1.72-1.95(5H,m), 2.53-2.65(1H,m), 5.04(1H,brs), 7.37(2H,d,J=8.2Hz), 7.81(2H,d,J=8.2Hz), 9.97(1H,s).

Example 1

4-(N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoic acid

(1) 1-(4-methoxycarbonylbenzyl)-2-thiourea


[0149]



[0150] Under an argon atmosphere, to a suspension of methyl 4-aminomethylbenzoate hydrochloride (170.0 g, 0.843 mol) in chloroform (850 ml, 5.0 v/w) were successively added 1,1'-thiocarbonyldiimidazole (purity 90%, 166.0 g, 0.843 mol) and triethylamine (123 ml, 0.885 mol). After stirring at room temperature for 3 hr, 28% aqueous ammonia (570 ml, 8.43 mol) and methanol (170 ml, 1.0 v/w) were added and the mixture was stirred overnight. n-Hexane (1700 ml, 10.0 v/w) and water (850 ml, 5.0 v/w) were successively added to the reaction mixture and the mixture was stirred at room temperature for 3 hrs. The precipitated crystals were collected by filtration, washed successively with n-hexane (500 ml) and water (500 ml) and dried in vacuo to give the title compound (172.5 g, yield 91.3%) as a colorless solid. ¹H-NMR(300MHz,DMSO-d₆) δ 3.84(3H,s), 4.40(1H,brs), 4.72 (2H,brs), 7.17(1H,brs), 7.41(2H,d,J=8.1Hz), 7.93(2H,d,J=8.4Hz), 8.07(1H,brs).

(2) methyl 4-((4-(4-nitrophenyl)-2-thiazolyl)aminomethyl)benzoate


[0151]



[0152] To a suspension of 1-(4-methoxycarbonylbenzyl)-2-thiourea (138.0 g, 0.554 mol) obtained in Example 1(1) in acetonitrile (1380 ml, 10.0 v/w) were successively added 2-bromo-4'-nitroacetophenone (124.1 g, 0.554 mol) and sodium bicarbonate (46.9 g, 0.559 mol), and the mixture was heated under reflux for 2 hr. After cooling to room temperature, water (1380 ml, 10.0 v/w) and n-hexane (690 ml, 5.0 v/w) were successively added and the mixture was stirred for 1 hr. The precipitated crystals were collected by filtration, washed successively with water (1000 ml) and n-hexane (500 ml) and dried in vacuo to give the title compound (183.9 g, yield 89.9%) as a yellow solid. ¹H-NMR(300MHz,DMSO-d₆) δ 3.83(3H,s), 4.63(2H,d,J=5.9Hz), 7.49(1H,s), 7.54(2H,d,J=8.1Hz), 7.95(2H,d,J=8.1Hz), 8.06(2H,d,J=9.3Hz), 8.23(2H,d,J=6.0Hz), 8.43(1H,t,J=5.9Hz).

(3) methyl 4-(N-methyl-N-(4-(4-nitrophenyl)-2-thiazolyl)aminomethyl)benzoate

[0153]




[0154] Under an argon atmosphere, to a suspension of sodium hydride (content 60%, 26.4 g, 0.661 mol) in N,N-dimethylformamide (530 ml, 2.5 v/w) were successively added dropwise a solution of methyl 4-((4-(4-nitrophenyl)-2-thiazolyl)aminomethyl)benzoate (212.4 g, 0.575 mol) obtained in Example 1(2) in N,N-dimethylformamide (743 ml, 3.5 v/w) and methyl iodide (41.2 ml, 0.661 mol) at 10°C or below, and the mixture was stirred at room temperature for 2 hr. Sodium hydride (content 60%, 2.3 g, 0.057 mol) was added and the mixture was stirred at room temperature for 1 hr. The reaction mixture was added dropwise to water (2120 ml, 10.0 v/w) at 10°C or below, and, after stirring at room temperature for 30 min., diisopropyl ether (848 ml, 4.0 v/w) was added and the mixture was stirred for 2 hr. The precipitated crystals were collected by filtration, washed successively with diisopropyl ether (424 ml) and water (424 ml) and dried in vacuo to give the title compound (201.9 g, yield 91.4%) as a yellow solid.

¹H-NMR(300MHz,DMSO-d₆) δ 3.14(3H,s), 3.84 (3H,s), 4.87(2H,s), 7.48(2H,d,J=8.3Hz), 7.61(1H,s), 7.96(2H,d,J=8.3Hz), 8.11(2H,d,J=9.0Hz), 8.25(2H,d,J=9.0Hz).

(4) methyl 4-(N-(4-(4-aminophenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate

[0155]




[0156] To a suspension of methyl 4-(N-methyl-N-(4-(4-nitrophenyl)-2-thiazolyl)aminomethyl)benzoate (200.0 g, 0.522 mol) obtained in Example 1(3) in a mixture of ethanol (800 ml, 4.0 v/w) and tetrahydrofuran (800 ml, 4.0 v/w) was added 10% palladium carbon (20.0 g) and the mixture was stirred overnight under a hydrogen atmosphere at 3 atm. The reaction mixture was filtered through celite and 10% palladium carbon (20.0 g) was added to the filtrate. The mixture was stirred under a hydrogen atmosphere at 3 atm for 3 hr. After celite filtration and solvent evaporation, toluene (800 ml) was added to the residue and the solvent was evaporated to give the title compound (182.5 g, yield 99.0%) as a yellow solid.

¹H-NMR(300MHz,DMSO-d₆) δ 3.07(3H,s), 3.84(3H,s), 4.82 (2H,s), 5.18(2H,br), 6.54(2H,d,J=8.6Hz), 6.78(1H,s), 7.45 (2H,d,J=8.3Hz), 7.52(2H,d,J=8.5Hz), 7.95(2H,d,J=8.2Hz).

(5) methyl 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate

[0157]




[0158] Under an argon stream, tetrahydrofuran (1000 ml, 5.7 v/w) was added to methyl 4-(N-(4-(4-aminophenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate (174.0 g, 0.493 mol) obtained in Example 1(4) and dissolved therein. A solution of 4-cyclohexylbenzaldehyde (120.6 g, 0.641 mol) obtained in Production Example 1(2) in tetrahydrofuran

(740 ml, 4.3 v/w) was added. Acetic acid (56.4 ml, 0.986 mol) was added and the mixture was stirred at room temperature for 1 hr. With stirring under ice-cooling, sodium triacetoxyborohydride (104.5 g, 0.493 mol) was added and the mixture was stirred at room temperature for 1.5 hr. After ice-cooling, acetic acid (28.2 ml, 0.493 mol) and sodium triacetoxyborohydride (52.2 g, 0.246 mol) were added and the mixture was stirred at room temperature for 1.5 hr. The reaction mixture was ice-cooled and carefully added to a saturated aqueous sodium hydrogen carbonate solution (2262 ml, 13.0 v/w). After stirring at room temperature for 1 hr, the mixture was extracted with ethyl acetate (522 ml), and the organic layer was washed successively with distilled water (174 ml) and saturated brine (522 ml) and dried over magnesium sulfate (50.0 g). The solvent was evaporated and the obtained orange solid was purified by silica gel column chromatography (developing solvent chloroform:ethyl acetate=99.5:0.5) to give the title compound (80.3 g, yield 31.0%) as a pale-yellow solid.

¹H-NMR(300MHz,DMSO-d₆) δ 1.16-1.47(5H,m), 1.65-1.80(5H,m), 2.40-2.55(1H,m), 3.60(3H,s), 3.83(3H,s), 4.22(2H,d,J=5.5Hz), 4.81(2H,s), 6.32(1H,t,J=5.5Hz), 6.57(1H,d,J=8.7Hz), 6.77(1H,s), 7.15(2H,d,J=8.1Hz), 7.26(2H,d,J=8.1Hz), 7.45(2H,d,J=8.3Hz), 7.53(2H,d,J=8.6Hz), 7.94(2H,d,J=8.3Hz).

(6) methyl 4-(N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate

[0159]




[0160] Under an argon stream, N,N-dimethylacetamide (351 ml, 5.0 v/w) was added to methyl 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate (70.3 g, 0.134 mol) obtained in Example 1(5) and potassium carbonate (73.9 g, 0.535 mol) was carefully added with stirring. After stirring the mixture at room temperature for 20 min., dimethyl sulfate (50.6 ml, 0.535 mol) was added. After stirring at 50°C for 1 hr., potassium carbonate (18.4 g, 0.134 mol) and dimethyl sulfate (12.7 ml, 0.134 mol) were added and the mixture was stirred at 60°C for 2 hr. After cooling to room temperature, n-hexane (422 ml, 6.0 v/w) was added, and the mixture was stirred for 1 hr. After ice-cooling, distilled water (562 ml, 8.0 v/w) was added. After stirring at room temperature for 1 hr., the resulting crystals were collected by filtration and washed with methanol (352 ml, 5.0 v/w) in a slurry form. Tetrahydrofuran (180 ml) was added to the obtained orange solid and insoluble materials were filtered off. The filtrate was concentrated to give the title compound (42.1 g, yield 58.3%) as a pale-yellow solid.

¹H-NMR(300MHz,DMSO-d₆) δ 1.16-1.47(5H,m), 1.65-1.80(5H,m), 2.40-2.55(1H,m), 3.01(3H,s), 3.07(3H,s), 3.83(3H,s), 4.54(2H,s), 4.81(2H,s), 6.71(2H,d,J=8.9Hz), 6.84(1H,s), 7.08-7.16(4H,m), 7.46(2H,d,J=8.3Hz), 7.62(2H,d,J=8.8Hz), 7.94(2H,d,J=8.3Hz).

(7) 4-(N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoic acid

[0161]



[0162] Under an argon stream, tetrahydrofuran (202 ml, 5.0 v/w) and methanol (102 ml, 3.0 v/w) were added to methyl 4-(N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate (40.7 g, 75.4 mmol) obtained in Example 1(6). With stirring at 50°C, a 1N-aqueous sodium hydroxide solution (151 ml, 151 mmol) was added. After stirring at 60°C for 1 hr., distilled water (173 ml, 4.25 v/w) was added. With stirring, 2N-hydrochloric acid (75.4 ml, 151 mmol) was carefully added. After stirring for 1 hr., the resulting crystals were collected by filtration and washed successively with distilled water (407 ml) and ethanol (204 ml), and dried in vacuo to give a yellow solid (39.2 g).

[0163] Tetrahydrofuran (172 ml, 4.5 v/w) was added to the obtained yellow solid (38.2 g) and the mixture was stirred at 50°C for 1 hr. After filtration, the residue was washed with tetrahydrofuran (19 ml, 0.5 v/w). Ethanol (134 ml, 3.5 v/w) and distilled water (134 ml, 3.5 v/w) were successively added to the filtrate with stirring at 50°C and the mixture was stirred at 50°C for 1 hr. and at room temperature for 1 hr. The resulting crystals were collected by filtration, washed with ethanol (306 ml) and dried in vacuo to give the title compound (36.5 g, yield 91.9%) as a pale-yellow solid.

¹H-NMR(300MHz,DMSO-d₆) δ 1.16-1.47 (5H,m), 1.65-1.80 (5H,m), 2.40-2.55(1H,m), 3.01(3H,s), 3.07(3H,s), 4.54(2H,s), 4.81(2H,s), 6.71(2H,d,J=9.0Hz), 6.83(1H,s), 7.08-7.16(4H,m), 7.43(2H,d,J=8.4Hz), 7.63(2H,d,J=8.8Hz), 7.92(2H,d,J=8.2Hz), 12.85(1H,brs).


melting point: 180-181°C

Example 2

potassium 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate

(1) 4-(4-cyclohexylbenzyloxy)acetophenone

[0164]




[0165] Toluene (225 ml) and 48% hydrobromic acid (150 ml) were added to 4-cyclohexylbenzyl alcohol (74.9 g, 0.394 mol) and the mixture was stirred at 50°C for 14 hr. After partitioning, the organic layer was washed successively with water (100 ml), saturated aqueous sodium hydrogen carbonate (100 ml), water (100 ml) and saturated brine, and dried over magnesium sulfate. After filtration and solvent evaporation, the residue was dried in vacuo to give a pale-yellow

oil. The obtained oil was dissolved in N,N-dimethylformamide (500 ml) and 4-hydroxyacetophenone (50.3 g, 0.369 mol) and potassium carbonate (153 g, 1.11 mol) were added. The mixture was stirred at 45°C for 70 min. After ice-cooling, water (750 ml) was added dropwise and the mixture was stirred at room temperature for 30 min. The precipitated crystals were collected by filtration, washed with water, washed with n-hexane and dried to give the title compound (93.5 g, yield 86.6%).

¹H-NMR(400MHz,DMSO-d₆) δ 1.15-1.52(5H,m), 1.58-1.87(5H,m), 2.47(1H,m), 2.51(3H,s), 5.15(2H,s), 7.10(2H,d, J=9.3Hz), 7.24(2H,d,J=8.4Hz), 7.36(2H,d,J=8.4Hz), 7.92(2H,d,J=9.3Hz).

(2) 2'-bromo-4-(4-cyclohexylbenzyloxy)acetophenone

[0166]




[0167] To a suspension of 4-(4-cyclohexylbenzyloxy)acetophenone (60.0 g, 0.195 mol) obtained in Example 2(1) in 1,2-dimethoxyethane (480 ml) was added dropwise a solution of bromine (10.5 ml, 0.205 mol) in 1,2-dimethoxyethane (120 ml) at room temperature. After stirring at room temperature for 90 min., water (600 ml) was added dropwise under ice-cooling, and the mixture was stirred at room temperature for 30 min. The precipitated crystals were collected by filtration, washed with water, washed with n-heptane and dried to give the title compound (68.2 g, yield 90.1%).

¹H-NMR(400MHz,CDCl₃) δ 1.18-1.50(5H,m), 1.70-1.95(5H,m), 2.51(1H,m), 4.38(2H,s), 5.10(2H,s), 7.03(2H,d, J=8.9Hz), 7.24(2H,d,J=8.4Hz), 7.34(2H,d,J=8.4Hz), 7.96(2H,d,J=8.9Hz).

(3) methyl 4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylaminomethyl)benzoate

[0168]




[0169] Acetonitrile (630 ml) was added to 1-(4-methoxycarbonylbenzyl)-2-thiourea (33.0 g, 0.147 mol) obtained in Example 1(1), 2'-bromo-4-(4-cyclohexylbenzyloxy)acetophenone (62.7 g, 0.162 mol) obtained in Example 2(2) and sodium bicarbonate (13.6 g, 0.162 mol), and the mixture was heated under reflux for 4 hr. After cooling to room temperature, water (630 ml) was added and the mixture was stirred at the same temperature for 1 hr. The precipitated crystals were collected by filtration, washed successively with 50% acetonitrile (130 ml), water (2 L) and diisopropyl

ether (500 ml) and dried to give the title compound (76.4 g, quant.).

¹H-NMR(400MHz,DMSO-d₆) δ 1.16-1.47(5H,m), 1.65-1.80(5H,m), 2.50(1H,m), 3.84(3H,s), 4.61(2H,brs), 5.06(2H,s), 6.91(1H,s), 7.00(2H,d,J=8.9Hz), 7.23(2H,d,J=8.1Hz), 7.35(2H,d,J=8.1Hz), 7.53(2H,d,J=8.1Hz), 7.70(2H,d,J=8.9Hz), 7.94(2H,d,J=8.1Hz), 8.28-8.52(1H,brs).

(4) methyl 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate


[0170]



[0171] Under an argon atmosphere, a suspension of methyl 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)benzoate (65.0 g, 0.127 mol) obtained in Example 2(3) in N,N-dimethylformamide (130 ml) and dimethyl sulfate (15.0 ml, 0.159 mol) were successively added dropwise to a suspension of sodium hydride (content 60%, 6.09 g, 0.152 mol) in N,N-dimethylformamide (130 ml) at 10°C or below and the mixture was stirred at room temperature for 1 hr. Diisopropyl ether (195 ml) and water (130 ml) were successively added dropwise at 10°C or below, and the mixture was stirred at room temperature for 30 min. The precipitated crystals were collected by filtration, washed successively with diisopropyl ether (195 ml) and water (130 ml) and dried to give the title compound (56.0 g, 83.7%).
¹H-NMR(400MHz,DMSO-d₆) δ 1.16-1.47(5H,m), 1.65-1.86(5H,m), 2-50(1H,m), 3.09(3H,s), 3.84(3H,s), 4.83(2H,s), 5.06(2H,s), 7.00(2H,d,J=8.9Hz), 7.01(1H,s), 7.23(2H,d,J=8.1Hz), 7.35(2H,d,J=8.1Hz), 7.46(2H,d,J=8.1Hz), 7.76(2H,d,J=8.9Hz), 7.94(2H,d,J=8.1Hz).

(5) 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoic acid

[0172]




[0173] Tetrahydrofuran (250 ml), methanol (250 ml) and a 2N-aqueous sodium hydroxide solution (95.0 ml, 190 mmol) were added to methyl 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate (50.0 g, 94.9 mmol) obtained in Example 2(4), and the mixture was heated under reflux for 40 min under an argon atmosphere. Water (310 ml) was added to the reaction mixture and the mixture was cooled to room temperature.

2N-Hydrochloric acid (95.0 ml, 190 mmol) was added dropwise and the mixture was stirred for 90 min. The resulting crystals were collected by filtration, washed with water (700 mL) and dried to give the title compound (48.5 g, yield 99.7%).

(6) potassium 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate

[0174]



[0175] Under an argon atmosphere, to a suspension of 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoic acid (30.0 g, 58.5 mmol) obtained in Example 2(5) was added 1N aqueous potassium hydroxide solution (56.0 ml) at 50°C and the mixture was heated under reflux for 40 min. After stirring at room temperature for 45 min., the crystals were collected by filtration, washed with a tetrahydrofuran-ethanol mixed solvent (3:1, 150 ml) and ethanol (210 ml), and dried to give the title compound (28.0 g, yield 90.9%).


¹H-NMR(400MHz,DMSO-d₆) δ 1.17-1.47(5H,m), 1.65-1.84(5H,m), 2.50(1H,m), 3.04(3H,s), 4.71(2H,s), 5.07(2H,s), 6.99(1H,s), 7.00(2H,d,J=8.9Hz), 7.19(2H,d,J=8.1Hz), 7.23(2H,d,J=8.1Hz), 7.36(2H,d,J=8.1Hz), 7.77(2H,d,J=8.9Hz), 7.79(2H,d,J=8.1Hz). melting point: 288-291°C (dec.)

Example 3

4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid

(1) ethyl 4-((4-(4-nitrophenyl)-2-thiazolyl)amino)benzoate hydrobromide

[0176]




[0177] A solution of 2'-bromo-4-nitroacetophenone (87.1 g, 0.357 mol) and 1-(4-ethoxycarbonylphenyl)-2-thiourea (80.0 g, 0.357 mol) in acetonitrile (1.6 L) was heated under reflux for 1 hr under an argon atmosphere. After cooling to room temperature, the resulting crystals were collected by filtration and dried to give the title compound (153 g, yield 94.9%).

¹H-NMR(400MHz,DMSO-d₆) δ 1.33(3H,t,J=7.1Hz), 4.29(2H,q,J=7.1Hz), 7.84(1H,s), 7.87(2H,d,J=9.1Hz), 7.97(2H,d,J=8.6Hz), 8.22(2H,d,J=8.6Hz), 8.27(2H,d,J=9.2Hz), 10.82(1H,s).

(2) ethyl 4-(N-methyl-N-(4-(4-nitrophenyl)-2-thiazolyl)amino)benzoate

[0178]




[0179] Under a nitrogen stream, N,N-dimethylformamide (1.05 L, 7.0 v/w) was added to ethyl 4-((4-(4-nitrophenyl)-2-thiazolyl)amino)benzoate hydrobromide (150 g, 0.333 mol) obtained in Example 3(1). With stirring under ice-cooling, potassium carbonate (138 g, 0.999 mol) was added carefully. After stirring at room temperature for 20 min., dimethyl sulfate (63.2 ml, 0.666 mol) was added. After stirring at 60°C for 2 hr., distilled water (1.05 L, 7.0 v/w) was added with stirring under ice-cooling. After stirring under ice-cooling for 1 hr., the resulting crystal were collected by filtration, washed with distilled water (450 mL) and dried in vacuo to give the title compound (127 g, yield 99.3%) as an orange crystal.

¹H-NMR(400MHz,DMSO-d₆) δ 1.34(3H,t,J=7.1Hz), 3.62(3H,s), 4.33(2H,q,J=7.1Hz), 7.74(2H,d,J=8.8Hz), 7.78(1H,s), 8.03(2H,d,J=8.8Hz), 8.14(2H,d,J=9.0Hz), 8.28(2H,d,J=9.0Hz).

(3) ethyl 4-(N-(4-(4-aminophenyl)-2-thiazolyl)-N-methylamino)benzoate

[0180]




[0181] Under a nitrogen stream, N,N-dimethylformamide (1.20 L, 10.0 v/w) was added to ethyl 4-(N-methyl-N-(4-(4-nitrophenyl)-2-thiazolyl)amino)benzoate (123 g, 0.321 mol) obtained in Example 3(2). With stirring at room temperature, sodium hydrosulfite (80% purity, 210 g, 0.963 mol) was added. After stirring at room temperature for 10 min., distilled water (123 ml, 1.0 v/w) was added carefully. After stirring at 100°C for 1.5 hr., triethylamine (223 ml, 1.61 mol) was added at 80°C, and the mixture was cooled to room temperature with water. After stirring at room temperature for 1 hr., distilled water (1.1 L, 9.0 v/w) was added. After stirring at room temperature for 30 min., the mixture was extracted twice with ethyl acetate (1.20 L) and the organic layer was washed successively with distilled water (400 ml) and saturated brine (400 ml), and dried over magnesium sulfate (60 g). The magnesium sulfate was filtered off, the solvent was evaporated, the residue was boiled with toluene and dried in vacuo to give the title compound (67.0 g, yield 63.0%) as a yellow to orange solid.

¹H-NMR(400MHz,DMSO-d₆) δ 1.33(3H,t,J=7.0Hz), 3.57(3H,s), 4.32(2H,q,J=7.0Hz), 5.27(2H,brs), 6.59(2H,d,J=8.6Hz), 7.02(1H,s), 7.56(2H,d,J=8.6Hz), 7.71(2H,d,J=9.0Hz), 7.99(2H,d,J=9.0Hz).

(4) ethyl 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoate

[0182]




[0183] Under a nitrogen stream, tetrahydrofuran (306 ml, 6.0 v/w) was added to ethyl 4-(N-(4-(4-aminophenyl)-2-thiazolyl)-N-methylamino)benzoate (51.0 g, 0.144 mol) obtained in Example 3(3). With stirring at room temperature, a solution of 4-cyclohexylbenzaldehyde (30.0 g, 0.158 mol) obtained in Production Example 1(2) in tetrahydrofuran (153 ml, 3.0 v/w) was added. The mixture was washed with tetrahydrofuran (51 ml, 1.0 v/w) and stirred at room temperature for 30 min. With stirring under ice-cooling, sodium triacetoxyborohydride (46.0 g, 0.216 mol) and acetic acid (12.4 ml, 0.216 mol) were added, and the mixture was stirred at room temperature for 1.5 hr. With stirring under ice-cooling, saturated aqueous sodium hydrogen carbonate (510 ml, 10.0 v/w) was added carefully. After stirring at room temperature for 1 hr., the mixture was extracted with ethyl acetate (408 ml) and the organic layer was washed successively with distilled water (255 ml) and saturated brine (255 ml), and dried over magnesium sulfate (50 g). After filtration and solvent evaporation, isopropyl alcohol (510 ml, 10.0 v/w) was added to the obtained orange solid, and the mixture was stirred at 60°C for 1 hr. After stirring under ice-cooling for 1 hr., the resulting crystal were collected by filtration, and washed with isopropyl alcohol (102 ml) and tert-butylmethyl ether (102 ml) and dried in vacuo to give the title compound (58.0 g, yield 77.0%) as a pale-yellow solid.

¹H-NMR(400MHz,DMSO-d₆) δ 1.23-1.40(8H,m), 1.67-1.78(5H,m), 2.46(1H,s), 3.56(3H,s), 4.24(2H,d,J=6.1Hz), 4.31(2H,q,J=7.1Hz), 6.35(1H,t,J=6.1Hz), 6.60(2H,d,J=8.6Hz), 7.01(1H,s), 7.16(2H,d,J=7.6Hz), 7.27(6H,d,J=7.6Hz), 7.57(2H,d,J=8.6Hz), 7.70(2H,d,J=8.9Hz), 7.98(2H,d,J=8.9Hz).

(5) 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid

[0184]



[0185] Under a nitrogen stream, tetrahydrofuran (312 ml, 6.0 v/w) and methanol (104 ml, 2.0 v/w) were added to ethyl 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoate (52.0 g, 98.9 mmol) obtained in Example 3(4). With stirring under ice-cooling, 2N-aqueous sodium hydroxide solution (98.9 ml, 197.8 mmol) was added. After stirring at 60°C for 2 hr., distilled water (104 ml, 2.0 v/w) was added. With stirring under ice-cooling, 2N-hydrochloric acid (98.9 ml, 197.8 mmol) was added carefully. After stirring under ice-cooling for 1 hr., the resulting crystal were collected by filtration, washed with distilled water (156 ml) and dried in vacuo to give a yellow solid (51.9 g). Tetrahydrofuran (750 ml, 15.0 v/w) was added to the obtained yellow solid (50.0 g) and the mixture was stirred at 60°C for 1 hr. After allowing to cool to room temperature, the precipitate was collected by filtration and washed with tetrahydrofuran (100 ml, 2.0 v/w). With stirring at room temperature, ethanol (150 ml) and distilled water (150 ml) were successively added to the filtrate. After stirring under ice-cooling for 1 hr., the resulting crystals were collected by filtration, washed successively with distilled water (200 ml) and 50% aqueous ethanol (200 ml) and dried in vacuo to

give a crude title compound (33.6 g) as a yellow solid. Ethanol (350 ml, 7.0 v/w) was added and the mixture was stirred at room temperature for 2 hr. The resulting crystals were collected by filtration, washed with ethanol (200 ml) and dried in vacuo to give the title compound (32.1 g, yield 64.2%) as a pale-yellow solid.

¹H-NMR(400MHz,DMSO-d₆) δ 1.20-1.40(5H,m), 1.67-1.78(5H,m), 2.43(1H,m), 3.56(1H,s), 4.23(3H,d,J=5.1Hz), 6.34(1H,brt,J=5.1Hz), 6.60(2H,d,J=8.6Hz), 6.98(1H,s), 7.16(2H,d,J=8.1Hz), 7.27(2H,d,J=8.1Hz), 7.57(2H,d,J=8.6Hz), 7.67(2H,d,J=8.9Hz), 7.97(2H,d,J=8.9Hz).

melting point: 252-253°C (dec.)

[0186] The following compounds were produced according to a method similar to the methods of Examples 1-3, and using a conventional method where necessary.

- 10 4-(4-(4-benzoylamino)phenyl)-2-thiazolylamino)benzoic acid (Example 4),
4-(4-(4-(4-tert-butylbenzoylamino)phenyl)-2-thiazolylamino)benzoic acid (Example 5),
4-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolylamino)benzoic acid (Example 6),
4-(N-(4-(4-benzoylamino)phenyl)-2-thiazolyl)-N-ethylamino)benzoic acid (Example 7),
4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-ethylamino)benzoic acid (Example 8),
15 4-(N-(4-(4-benzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 9),
4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 10),
4-(N-(4-(4-(4-cyclohexanecarbonylamino)phenyl)-2-thiazolyl)-N-ethylamino)benzoic acid (Example 11),
4-(N-(4-(4-(4-cyclohexanecarbonylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 12),
4-(N-(4-(4-(4-cyclohexanecarbonylamino)phenyl)-2-thiazolyl)-N-isobutylamino)benzoic acid (Example 13),
20 4-(N-carboxymethyl-N-(4-(4-(4-cyclohexanecarbonylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 14),
4-(N-(4-(4-benzoylamino)phenyl)-2-thiazolyl)-N-isobutylamino)benzoic acid (Example 15),
4-(N-(4-(4-benzoylamino)phenyl)-2-thiazolyl)-N-carboxymethylamino)benzoic acid (Example 16),
4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isobutylamino)benzoic acid (Example 17),
4-(N-carboxymethyl-N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 18),
25 4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 19),
4-(N-(4-(4-(4-tert-butylbenzoylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 20),
4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 21),
4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 22),
4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-ethylamino)benzoic acid (Example 23),
30 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 24),
4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-cyclohexylmethylamino)benzoic acid (Example 25),
4-(N-(3-carboxypropyl)-N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)amino)benzoic acid (Example 26),
3-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolylamino)benzoic acid (Example 27),
3-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 28),
35 4-(N-isopropyl-N-(4-(4-(4-morpholinobenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 29),
3-(N-isopropyl-N-(4-(4-(4-piperidinobenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 30),
3-(N-isopropyl-N-(4-(4-(4-morpholinobenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 31),
40 4-(N-isopropyl-N-(4-(4-(4-piperidinobenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 32),
4-(N-isopropyl-N-(4-(4-(4-methylpiperidino)benzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 33),
45 sodium 4-(N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino) benzoate (Example 34),
4-(N-(4-(4-(4-(3,5-dimethylpiperidino)benzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid hydrochloride (Example 35),
cis-4-(N-(4-(4-(4-(2,6-dimethylmorpholino)benzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid hydrochloride (Example 36),
50 sodium 4-(N-isopropyl-N-(4-(4-(4-methyl-1-piperazinyl)benzoylamino)phenyl)-2-thiazolyl)amino)benzoate (Example 37),
2-chloro-4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 38),
2-chloro-4-(N-isopropyl-N-(4-(4-(4-piperidinobenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 39),
55 2-chloro-4-(N-isopropyl-N-(4-(4-(4-morpholinobenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 40),
2-chloro-4-(N-isopropyl-N-(4-(4-(4-methyl-1-piperazinyl)benzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 41),

- 4-(1-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-1-methylethyl)benzoic acid (Example 42),
 4-(1-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-1-methylethyl)benzoic acid (Example 43),
 4-(1-methyl-1-(4-(4-(4-morpholinobenzoylamino)phenyl)-2-thiazolyl)ethyl)benzoic acid (Example 44),
 4-(1-methyl-1-(4-(4-(4-piperidinobenzoylamino)phenyl)-2-thiazolyl)ethyl)benzoic acid (Example 45),
 5 4-(N-(4-(4-(4-cyclohexylbutyrylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 46),
 4-(4-(4-(4-tert-butylbenzyloxy)phenyl)-2-thiazolylmethyl)benzoic acid (Example 47),
 4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylmethyl)benzoic acid (Example 48),
 4-(4-(4-(4-carboxybenzyloxy)phenyl)-2-thiazolylmethyl)benzoic acid (Example 49),
 (4-(N-(4-(4-(cyclohexanecarbonylamino)phenyl)-2-thiazolyl)-N-isopropylamino)phenoxy)acetic acid (Example 50),
 10 (4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)phenoxy)acetic acid (Example 51),
 4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)-2,3,5,6-tetrafluorobenzoic acid (Example 52),
 (4-(N-isopropyl-N-(4-(4-(4-piperidinobenzoylamino)phenyl)-2-thiazolyl)amino)phenoxy)acetic acid (Example 53),
 (4-(N-isopropyl-N-(4-(4-(4-morpholinobenzoylamino)phenyl)-2-thiazolyl)amino)phenoxy)acetic acid (Example 54),
 15 (4-(N-(4-(4-(3,5-bis(trifluoromethyl)benzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)phenoxy)acetic acid (Example 55),
 (4-(N-(4-(4-(3,5-dichlorobenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)phenoxy)acetic acid (Example 56),
 (4-(N-isopropyl-N-(4-(4-(2-piperidino-5-pyridinecarbonylamino)phenyl)-2-thiazolyl)amino)phenoxy)acetic acid (Example 57),
 20 4-(N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-cyclohexylmethylamino)benzoic acid (Example 58),
 4-(N-cyclohexylmethyl-N-(4-(4-(4-trifluoromethylbenzoylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 59),
 4-(N-(4-(4-(N-(4-cyclohexylbenzoyl)-N-methylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 60),
 4-((N-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 61),
 25 4-((N-isopropyl-N-(4-(4-(4-piperidinobenzoylamino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 62),
 4-((N-isopropyl-N-(4-(4-(4-morpholinobenzoylamino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 63),
 4-((N-(4-(4-(4-biphenylcarbonylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 64),
 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 65),
 4-(2-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)ethyl)benzoic acid (Example 66),
 30 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 67),
 4-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolylmethyl)benzoic acid (Example 68),
 4-(4-(4-(4-isopropoxybenzoylamino)phenyl)-2-thiazolylmethyl)benzoic acid (Example 69),
 4-(4-(4-(4-(1-pyrrolyl)benzoylamino)phenyl)-2-thiazolylmethyl)benzoic acid (Example 70),
 4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylmethyl)benzoic acid (Example 71),
 35 4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylamino)-3-(2-cyclohexylethoxy)benzoic acid (Example 72),
 3-benzyloxy-4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 73),
 3-(4-carboxybenzyloxy)-4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 74),
 4-cyclohexyl-N-(4-(2-(N-isopropyl-N-(4-(1H-tetrazol-5-yl)phenyl)amino)-4-thiazolyl)phenyl)benzamide (Example 75),
 3-(2-(4-(4-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)ethyl)benzoic acid (Example 76),
 40 4-((N-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 77),
 N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropyl-(4-(1H-tetrazol-5-yl)phenyl)amine (Example 78),
 N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropyl-(4-(1H-tetrazol-5-yl)benzyl)amine (Example 79),
 N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropyl-(3-(1H-tetrazol-5-yl)benzyl)amine (Example 80),
 4-((4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylamino)methyl)benzoic acid (Example 81),
 45 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 82),
 4-((N-(4-(4-(4-fluorophenyl)-2-methyl-5-thiazolylmethoxy)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 83),
 4-((N-(4-(4-(4'-chloro-4-methoxybiphenyl-2-ylmethoxy)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 84),
 50 4-((N-isopropyl-N-(4-(4-(4-methyl-2-(4-trifluoromethylphenyl)-5-thiazolylmethoxy)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 85),
 N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropyl-(2-(1H-tetrazol-5-yl)benzyl)amine (Example 86),
 3-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 87),
 2-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 88),
 55 4-(N-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 89),
 4-(1-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)-1-methylethyl)benzoic acid (Example 90),
 4-(1-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)cyclopentyl)benzoic acid (Example 91),
 4-(1-(4-(4-(4-biphenylmethoxy)phenyl)-2-thiazolyl)-1-methylethyl)benzoic acid (Example 92),

- 4-(1-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)cyclohexyl)benzoic acid (Example 93),
 4-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 94),
 4-(4-(4-(4-biphenylmethoxy)phenyl)-2-thiazolylamino)benzoic acid (Example 95),
 4-(N-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 96),
 5 4-(N-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)-N-(2-dimethylaminoethyl)amino)benzoic acid (Example 97),
 4-(N-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)-N-(2-piperidinoethyl)amino)benzoic acid (Example 98),
 4-(N-(4-(4-(3,4-dichlorobenzyloxy)phenyl)-2-thiazolyl)-N-(2-methoxyethyl)amino)benzoic acid (Example 99),
 4-(1-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)cyclohexyl)benzoic acid (Example 100),
 4-(4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)tetrahydropyran-4-yl)benzoic acid (Example 101),
 10 sodium 4-(N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylaminomethyl)benzoate (Example 102),
 4-((N-isopropyl-N-(4-(4-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 103),
 4-((N-(4-(4-(4-tert-butylbenzenesulfonylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 104),
 15 4-((N-(4-(4-(3,4-dichlorobenzenesulfonylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 105),
 4-((N-isopropyl-N-(4-(4-(4-trifluoromethylbenzenesulfonylamino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 106),
 4-((N-(4-(4-(4-cyclohexylbenzenesulfonylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 107),
 20 4-((N-(4-(4-(4-cyclohexylbenzylsulfanyl)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 108),
 4-((N-(4-(4-dibenzylaminophenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 109),
 4-((N-(4-(4-(N-(4-cyclohexylbenzenesulfonyl)-N-methylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 110),
 25 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-propylamino)methyl)benzoic acid (Example 111),
 4-((N-(4-(4-(4-cyclohexylphenylmethanesulfonyl)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 112),
 4-((N-benzyl-N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 113),
 4-((N-(4-(4-(N-benzyl-N-(4-cyclohexylbenzyl)amino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 114),
 30 4-((N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 115),
 4-((N-cyclohexylmethyl-N-(4-(4-(3,4-dichlorobenzylamino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 116),
 4-((N-(4-(4-(bis(3,4-dichlorobenzyl)amino)phenyl)-2-thiazolyl)-N-cyclohexylmethylamino)methyl)benzoic acid (Example 117),
 35 4-((N-cyclohexylmethyl-N-(4-(4-(N-(3,4-dichlorobenzyl)-N-methylamino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 118),
 4-((N-(4-(4-(N-allyl-N-(4-cyclohexylbenzyl)amino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 119),
 40 sodium 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 120),
 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid N-methyl-D-glucamine salt (Example 121),
 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid tris(hydroxymethyl)aminomethane salt (Example 122),
 45 4-((N-(4-(4-(N-benzyl-N-cyclohexylmethylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 123),
 4-((N-(4-(4-(N-cyclohexylmethyl-N-(4-trifluoromethylbenzyl)amino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 124),
 sodium 4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolecarbonyl)benzoate (Example 125),
 50 potassium 4-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolecarbonyl)benzoate (Example 126),
 4-((N-isopropyl-N-(4-(4-(3-piperidinobenzyloxy)phenyl)-2-thiazolyl)amino)methyl)benzoic acid dihydrochloride (Example 127),
 4-(N-(4-(4-(3,4-dichlorobenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 128),
 4-(N-cyclohexylmethyl-N-(4-(4-(3,4-dichlorobenzylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 129),
 55 4-(N-cyclohexylmethyl-N-(4-(4-(4-isopropylbenzylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 130),
 4-(N-cyclohexylmethyl-N-(4-(4-(4-isobutylbenzylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 131),
 4-(N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 132),
 4-(N-methyl-N-(4-(4-(4-trifluoromethylbenzylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 133),

- 4-(N-cyclohexylmethyl-N-(4-(4-(4-trifluoromethylbenzylamino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 134),
 4-((N-(4-(4-(N-cyclohexylmethyl-N-(4-trifluoromethylbenzyl)amino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 135),
 4-((N-isopropyl-N-(4-(4-(N-(tetrahydropyran-4-ylmethyl)-N-(4-trifluoromethylbenzyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 136),
 4-(N-methyl-N-(4-(4-(N-methyl-N-(4-trifluoromethylbenzyl)amino)phenyl)-2-thiazolyl)amino)benzoic acid (Example 137),
 4-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 138),
 4-(N-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 139),
 4-(N-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-ethylamino)benzoic acid (Example 140),
 4-(N-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 141),
 4-(N-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-cyclohexylmethylamino)benzoic acid (Example 142),
 4-((N-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 143),
 4-((N-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 144),
 4-((N-(4-(3-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-cyclohexylmethylamino)methyl)benzoic acid (Example 145),
 4-((N-(4-(3-(4-cyclohexylbenzoylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 146),
 4-((N-(4-(3-(4-cyclohexylbenzenesulfonylamino)phenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 147),
 4-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolylamino)benzoic acid (Example 148),
 4-(1-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-1-methylethyl)benzoic acid (Example 149),
 4-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolylamino)-3-(2-cyclohexylethoxy)benzoic acid (Example 150),
 4-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-methylamino)benzoic acid (Example 151),
 4-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 152),
 4-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-(2-phenoxyethyl)amino)benzoic acid (Example 153),
 4-(4-(2-benzyloxy-5-methoxyphenyl)-2-thiazolylamino)benzoic acid (Example 154),
 4-(4-(5-methoxy-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 155),
 3-benzyloxy-4-(4-(2-benzyloxy-5-methoxyphenyl)-2-thiazolylamino)benzoic acid (Example 156),
 4-(4-(2-(4-cyclohexylbenzyloxy)-5-methoxyphenyl)-2-thiazolylamino)benzoic acid (Example 157),
 5-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolylmethyl)-2-hydroxybenzoic acid (Example 158),
 4-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-(4-trifluoromethylbenzyl)amino)benzoic acid (Example 159),
 4-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-cyclohexylmethylamino)benzoic acid (Example 160),
 4-((N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 161),
 4-((N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolylamino)methyl)benzoic acid (Example 162),
 4-((N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 163),
 4-((N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)methyl)benzoic acid (Example 164),
 4-((N-(4-(5-chloro-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)methyl)benzoic acid (Example 165),
 4-((N-(2-cyclohexylethyl)-N-(4-(5-methoxy-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 166),
 4-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-(3,4-dichlorobenzyl)amino)benzoic acid (Example 167),
 4-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-(3-methylbutyl)amino)benzoic acid (Example 168),
 4-(N-(4-(2-benzyloxy-5-methoxyphenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 169),
 4-(N-(2-cyclohexylethyl)-N-(4-(5-methoxy-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolyl)amino)benzoic acid (Example 170),
 3-(2-cyclohexylethoxy)-4-(4-(5-methoxy-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 171),
 3-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 172),
 4-(4-(5-chloro-2-(4-isopropylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 173),
 4-(N-(4-(2-benzyloxy-5-fluorophenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 174),
 4-(N-(4-(2-benzyloxy-5-methylphenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 175),
 4-(N-(4-(5-chloro-2-(4-isopropylbenzyloxy)phenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 176),
 4-(N-(4-(5-chloro-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 177),
 4-(N-(4-(5-chloro-2-(3,4-dichlorobenzoyloxy)phenyl)-2-thiazolyl)-N-(2-cyclohexylethyl)amino)benzoic acid (Example 178),
 4-(4-(5-chloro-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolylamino)-3-(2-cyclohexylethoxy)benzoic acid (Example 179),

- 3-benzyloxy-4-(4-(5-methoxy-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolylamino)benzoic acid (Example 180),
 3-(N-(4-(2-benzyloxy-5-chlorophenyl)-2-thiazolyl)-N-cyclohexylmethylamino)benzoic acid (Example 181),
 4-(N-butyl-N-(4-(5-chloro-2-(4-trifluoromethylbenzyloxy)phenyl)-2-thiazolyl)amino)benzoic acid (Example 182),
 4-(N-(4-(6-(3,4-dichlorobenzyloxy)-2-benzooxazolyl)-2-thiazolyl)-N-isopropylamino)benzoic acid (Example 183),
 5 4-(N-(2-cyclohexylethyl)-N-(4-(6-(3,4-dichlorobenzyloxy)-2-benzooxazolyl)-2-thiazolyl)amino)benzoic acid (Example 184),
 4-(N-(2-cyclohexylethyl)-N-(4-(5-(3,4-dichlorobenzyloxy)-2-benzimidazolyl)-2-thiazolyl)amino)benzoic acid hydrochloride (Example 185),
 4-(4-(5-(3,4-dichlorobenzyloxy)-2-benzimidazolyl)-2-thiazolylamino)benzoic acid hydrochloride (Example 186),
 10 4-(N-cyclohexylmethyl-N-(4-(6-(3,4-dichlorobenzyloxy)-2-benzooxazolyl)-2-thiazolyl)amino)benzoic acid (Example 187),
 4-(N-(4-(6-(4-tert-butylbenzyloxy)-2-benzooxazolyl)-2-thiazolyl)-N-cyclohexylmethylamino)benzoic acid (Example 188),
 4-(N-(4-(6-(2,4-bis(trifluoromethyl)benzyloxy)-2-benzooxazolyl)-2-thiazolyl)-N-cyclohexylmethylamino)benzoic acid (Example 189),
 15 4-(N-cyclohexylmethyl-N-(4-(6-(3,4-difluorobenzyloxy)-2-benzooxazolyl)-2-thiazolyl)amino)benzoic acid (Example 190),
 4-(N-cyclohexylmethyl-N-(4-(6-(3,5-dimethoxybenzyloxy)-2-benzooxazolyl)-2-thiazolyl)amino)benzoic acid (Example 191),
 4-((N-(4-(4-cyclopentylloxyphenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid (Example 192),
 20 4-((N-(4-(4-dicyclohexylmethoxyphenyl)-2-thiazolyl)-N-isopropylamino)methyl)benzoic acid hydrochloride (Example 193),
 cis-4-((N-(4-(4-(N-(4-(4-hydroxycyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 194),
 4-((N-methyl-N-(4-(4-(N-methyl-N-(4-(4-oxocyclohexyl)benzyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 195),
 25 trans-4-((N-(4-(4-(N-(4-(4-hydroxycyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 196),
 4-((N-(4-(3-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 197),
 4-((N-(4-(4-(N-(4-(4-cyclohexylbenzyl)-N-(2-methoxyacetyl)amino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 198),
 30 4-((N-(4-(4-(N-(4-(4-cyclohexylbenzyl)-N-(2-hydroxyacetyl)amino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 199),
 sodium 4-((N-(4-(3-(N-(4-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 200),
 35 4-((N-(4-(3-(N-acetyl-N-(4-(4-cyclohexylbenzyl)amino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 201),
 4-((N-(4-(4-(N-(4-(4-cyclohexylbenzyl)-N-methanesulfonylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 202),
 4-((N-(4-(4-(N-(4-(4-cyclohexylbenzyl)-N-(2-hydroxyethyl)amino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 203),
 40 3-((N-(4-(4-(N-(4-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 204),
 cis-4-((N-(4-(4-(N-(4-(3-hydroxycyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 205),
 trans-4-((N-(4-(4-(N-(4-(3-hydroxycyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 206),
 45 4-((N-methyl-N-(4-(4-(N-methyl-N-(4-(3-oxocyclohexyl)benzyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 207),
 4-((N-(4-(4-(N-(4-(4,4-dichlorocyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 208),
 4-((N-(4-(4-(N-(4-(4,4-difluorocyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 209),
 50 4-((N-methyl-N-(4-(4-(N-methyl-N-(4-(tetrahydropyran-4-yl)benzyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 210),
 4-((N-(4-(4-(N-(4-(1-acetyl piperidin-4-yl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 211),
 55 4-((N-(4-(4-(N-(4-(4-cyclopentylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 212),
 trans-4-((N-methyl-N-(4-(4-(N-methyl-N-(4-(phenylcyclohexylmethyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 213),

- cis-4-((N-methyl-N-(4-(4-(N-methyl-N-(4-phenylcyclohexylmethyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 214),
- 4-((N-(4-(4-(N-(4-(4,4-dimethylcyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 215),
- 5 sodium 4-((N-(4-(4-(N-(4-(4,4-dimethylcyclohexyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 216),
- 4-((N-cyclohexylmethyl-N-(4-(6-(3,4-dichlorobenzyloxy)-2-benzooxazolyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 217),
- 4-((N-methyl-N-(4-(4-(N-methyl-N-(4-(4-methylcyclohexyl)benzyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 218),
- 10 4-((N-(4-(4-(N-(2-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 219), 4-((N-(4-(6-benzyloxy-2-benzooxazolyl)-2-thiazolyl)-N-cyclohexylmethylamino)methyl)benzoic acid (Example 220),
- 4-((N-cyclohexylmethyl-N-(4-(6-phenethyloxy-2-benzooxazolyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 221),
- 15 4-((N-methyl-N-(4-(4-(N-methyl-N-(4-phenylbutyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 222),
- 4-((N-(4-(4-(N-(2-(2-indanyl)ethyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 223),
- 20 4-((N-(4-(4-(N-(2-(3,4-dichlorophenyl)ethyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 224),
- 4-((N-methyl-N-(4-(4-(N-methyl-N-(3-phenylpropyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 225),
- 4-((N-(4-(6-(N-(4-cyclohexylbenzyl)-N-methylamino)-3-pyridyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 226),
- 25 4-((N-(2'-(N-(4-cyclohexylbenzyl)-N-methylamino)-4'-methyl-4,5'-bithiazolyl-2-yl)-N-methylamino)methyl)benzoic acid (Example 227),
- 4-((N-(4-(2-chloro-6-(N-(4-cyclohexylbenzyl)-N-methylamino)-4-pyridyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 228),
- 30 4-((N-(4-(4-(N-(4-(2,2-dimethylpropyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 229),
- 4-((N-methyl-N-(4-(4-(N-methyl-N-(4-(1-propylbutyl)benzyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 230),
- 4-((N-(4-(2-(4-cyclohexylphenyl)-1H-benzimidazol-5-yl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 231),
- 35 4-((N-(4-(5-chloro-6-(N-(4-cyclohexylbenzyl)-N-methylamino)-3-pyridyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 232),
- 4-((N-(4-(4-(N-(4-tert-butylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 233),
- 40 4-((N-methyl-N-(4-(4-(N-methyl-N-(4-trifluoromethylbenzyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 234),
- 4-((N-(4-(4-(N-(4-tert-butylphenyl)ethyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 235),
- 4-((N-methyl-N-(4-(4-(N-methyl-N-(2-(4-trifluoromethylphenyl)ethyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 236),
- 45 4-((N-(4-(4-(N-(2-(4-dimethylaminophenyl)ethyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 237),
- 4-((N-methyl-N-(4-(4-(N-methyl-N-(2-(4-morpholinophenyl)ethyl)amino)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 238),
- 50 4-((N-methyl-N-(4-(4-(4-phenyl-1-piperazinylmethyl)phenyl)-2-thiazolyl)amino)methyl)benzoic acid (Example 239),
- 4-((N-(4-(4-(4-benzyl-1-piperazinylmethyl)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 240),
- 4-((N-(4-(4-(N-(4-(1-ethylpropyl)benzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 241),
- 4-((N-(4-(3-chloro-4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 242),
- 55 benzyl 4-(4-(2-(N-(4-carboxybenzyl)-N-methylamino)-4-thiazolyl)benzyl)-piperazine-1-carboxylate (Example 243),
- 4-((N-(4-(4-(N-(4-tert-butylbenzyl)-N-methylamino)-3-chlorophenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 244),

- 4-((N-(4-(4-(N-(4-isobutylsulfanylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid (Example 245),
- 4-((N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid lysine salt (Example 246),
- 5 4-((4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolylamino)methyl)benzoic acid (Example 247),
- 3-oxo-1,3-dihydroisobenzofuran-1-yl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 248),
- ethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 249),
- 10 1-acetoxyethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 250),
- 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid methanesulfonate (Example 251),
- 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid hydrochloride (Example 252),
- 15 potassium 4-((N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 253),
- 4-((N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid hydrochloride (Example 254),
- 4-((N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid methanesulfonate (Example 255),
- 20 4-((N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid tris(hydroxymethyl)aminomethane salt (Example 256),
- 4-((N-(4-(4-(N-(4-cyclohexylbenzyl)-N-methylamino)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid sulfate (Example 257),
- 25 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid hydrochloride (Example 258),
- 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid methanesulfonate (Example 259),
- 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid tris(hydroxymethyl)aminomethane salt (Example 260),
- 30 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid N-methyl-D-glucamine salt (Example 261),
- 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid sulfate (Example 262),
- sodium 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoate (Example 263),
- 35 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid 1/2 sulfate (Example 264),
- 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid histidine salt (Example 265),
- 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoic acid lysine salt (Example 266),
- 40 tert-butoxycarbonylmethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 267),
- 2,2-dimethylpropionyloxymethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 268),
- 1-ethoxycarbonyloxyethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 269),
- 45 acetoxymethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 270),
- 1-isopropoxycarbonyloxyethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 271),
- 50 1-cyclohexyloxycarbonyloxyethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 272),
- 5-methyl-2-oxo-1,3-dioxl-4-ylmethyl 4-((N-(4-(4-(4-cyclohexylbenzyloxy)phenyl)-2-thiazolyl)-N-methylamino)methyl)benzoate (Example 273),
- 4-(N-(4-(4-(4-cyclohexylbenzylamino)phenyl)-2-thiazolyl)-N-methylamino)benzoic acid lysine salt (Example 274).
- 55 **[0187]** The structural formulas and property values of the compound of each Example are shown in Tables 1 to 67.

Table 1







Ex.	Structural formula	m.p. (°C)
1		180~ 181
2		288~ 291 (dec.)
3		252~ 253
4		250 (dec.)
5		250 (dec.)
6		250 (dec.)

Table 2

Ex.	Structural formula	m.p. (°C)
7		250 (dec.)
8		250 (dec.)
9		234- 236

Table 3




Ex.	Structural formula	m.p. (°C)
10		230 (dec.)
11		250 (dec.)
12		230 (dec.)

Table 4




Ex.	Structural formula	m.p. (°C)
13		231- 232
14		210 (dec.)
15		234- 235

Table 5





Ex.	Structural formula	m.p. (°C)
16		210 (dec.)
17		230 (dec.)
18		230 (dec.)
19		250 (dec.)

Table 6





Ex.	Structural formula	m.p. (°C)
20		250 (dec.)
21		250 (dec.)
22		250 (dec.)
23		239- 247

Table 7



Ex.	Structural formula	m.p. (°C)
24	 <chem>CC1CN(C1c2ccc(cc2)C(=O)O)c3nc(cc3c4ccc(cc4)Oc5ccc(cc5)C6CCCCC6)c5</chem>	238- 240
25	 <chem>C1CCN(C1Cc2ccc(cc2)C(=O)O)c3nc(cc3c4ccc(cc4)Oc5ccc(cc5)C6CCCCC6)c5</chem>	230 (dec.)

Table 8




Ex.	Structural formula	m.p. (°C)
26		230 (dec.)
27		230 (dec.)
28		230 (dec.)

Table 9




Ex.	Structural formula	m.p. (°C)
29	 <chem>CC(C)N1C(=N)SC1C2=CC=C(C=C2)C(=O)O</chem> <chem>ClH</chem>	230 (dec.)
30	 <chem>CC(C)N1C(=N)SC1C2=CC=C(C=C2)C(=O)O</chem> <chem>ClH</chem>	220 (dec.)
31	 <chem>CC(C)N1C(=N)SC1C2=CC=C(C=C2)C(=O)O</chem> <chem>ClH</chem>	230 (dec.)

Table 10




Ex.	Structural formula	m.p. (°C)
32	 <p>ClH</p>	230 (dec.)
33	 <p>ClH</p>	230 (dec.)
34	 <p>Na⁺</p>	230 (dec.)

Table 11



Ex.	Structural formula	m.p. (°C)
35	 <chem>CC(C)N(c1ccc(cc1)C(=O)O)c2nc(c3ccc(cc3NC(=O)c4ccc(cc4)N5CC(C)CC5)cc4)cs2.Cl</chem> ClH	152- 160
36	 <chem>CC(C)N(c1ccc(cc1)C(=O)O)c2nc(c3ccc(cc3NC(=O)c4ccc(cc4)N5CC(C)OC5)cc4)cs2.Cl</chem> ClH	230 (dec.)

Table 12




Ex.	Structural formula	m.p. (°C)
37		230 (dec.)
38		220 (dec.)
39		220 (dec.)

Table 14







Ex.	Structural formula	m.p. (°C)
44		290 (dec.)
45		268- 271
46		230 (dec.)
47		223- 228
48		235- 238
49		270 (dec.)

Table 15






Ex.	Structural formula	m.p. (°C)
50		225 (dec.)
51		240 (dec.)
52		153- 157
53		220 (dec.)
54		250 (dec.)

Table 16

Ex.	Structural formula	m.p. (°C)
55		amorphous
56		192-194
57		237-240
58		amorphous
59		amorphous

Table 17






Ex.	Structural formula	m.p. (°C)
60		amorphous
61		250 (dec.)
62		234 (dec.)
63		250 (dec.)
64		250 (dec.)

Table 18







Ex.	Structural formula	m.p. (°C)
65		233-234
66		amorphous
67		amorphous
68		230 (dec.)
69		230 (dec.)
70		220 (dec.)

Table 19






Ex.	Structural formula	m.p. (°C)
71		259-261
72		230 (dec.)
73		230 (dec.)
74		230 (dec.)
75		amorphous

Table 20

Ex.	Structural formula	m.p. (°C)
76		amorphous
77		180-181
78		215-216
79		238-239
80		221-222
81		amorphous

Table 21





Ex.	Structural formula	m.p. (°C)
82		amorphous
83		amorphous
84		amorphous
85		amorphous

Table 22







Ex.	Structural formula	m.p. (°C)
86		amorphous
87		196-197
88		259-261
89		230 (dec.)
90		170-172
91		248-251

Table 23







Ex.	Structural formula	m.p. (°C)
92		230 (dec.)
93		168- 171
94		230 (dec.)
95		230 (dec.)
96		236- 238
97		230 (dec.)

Table 24






Ex.	Structural formula	m.p. (°C)
98		230 (dec.)
99		175-178
100		222-224
101		230 (dec.)
102		240

Table 25






Ex.	Structural formula	m.p. (°C)
103		amorphous
104		178-180
105		193-195
106		132-134
107		amorphous

Table 26






Ex.	Structural formula	m.p. (°C)
108		amorphous
109		144-146
110		amorphous
111		amorphous
112		amorphous

Table 27






Ex.	Structural formula	m.p. (°C)
113		amorphous
114		amorphous
115		184 (dec.)
116		amorphous
117		120 (dec.)

Table 28






Ex.	Structural formula	m.p. (°C)
118		125-127
119		139-141
120		250 (dec.)
121		170-174
122		186-189

Table 29






Ex.	Structural formula	m.p. (°C)
123		amorphous
124		amorphous
125		270 (dec.)
126		270 (dec.)
127	 <p data-bbox="678 1717 737 1749">2HCl</p>	135-140

Table 30

Ex.	Structural formula	m.p. (°C)
128		223- 225
129		187- 189
130		200- 202
131		167- 169
132		185- 189

Table 31





Ex.	Structural formula	m.p. (°C)
133		230 (dec.)
134		199-203
135		amorphous
136		amorphous

Table 32





Ex.	Structural formula	m.p. (°C)
137		218-220
138		235-236
139		250 (dec.)
140		179-180

Table 33





Ex.	Structural formula	m.p. (°C)
141		189-190
142		195-197
143		amorphous
144		amorphous

Table 34





Ex.	Structural formula	m.p. (°C)
145		amorphous
146		228-230
147		amorphous
148		208-211

Table 35

Ex.	Structural formula	m.p. (°C)
149	 <chem>CC(C)(c1ccc(cc1)C(=O)O)c2nc(c(s2)Cc3ccc(Cl)cc3)Cc4ccccc4</chem>	152-154
150	 <chem>CC1(CCC1)COc2ccc(cc2Nc3nc(c(s3)Cc4ccc(Cl)cc4)Cc5ccccc5)C(=O)O</chem>	230 (dec.)
151	 <chem>CCN(C)c1nc(c(s1)Cc2ccc(Cl)cc2)Cc3ccccc3C(=O)O</chem>	211-212

Table 36




Ex.	Structural formula	m.p. (°C)
152		160- 163
153		152- 155
154		184- 185

Table 37

Ex.	Structural formula	m.p. (°C)
155	 <chem>COc1ccc(cc1C2=NC(=S)Nc3ccc(cc3)C(=O)O)COC4=CC=C(C(F)(F)F)C4</chem>	218- 219
156	 <chem>COc1ccc(cc1C2=NC(=S)Nc3ccc(cc3)C(=O)O)COC4=CC=C(COC5=CC=CC=C5)C4</chem>	218- 221
157	 <chem>COc1ccc(cc1C2=NC(=S)Nc3ccc(cc3)C(=O)O)COC4=CC=C(CCC5=CCCCC5)C4</chem>	230- 234

Table 38




Ex.	Structural formula	m.p. (°C)
158		212- 213
159		194- 196
160		184- 185

Table 39

Ex.	Structural formula	m.p. (°C)
161	 <chem>CC(C)N(Cc1ccc(cc1)C(=O)O)c2nc(c(s2)c3cc(Cl)ccc3OCc4ccccc4)cc5ccccc5</chem>	amorphous
162	 <chem>O=C(O)c1ccc(cc1)NCc2nc(c(s2)c3cc(Cl)ccc3OCc4ccccc4)cc5ccccc5</chem>	210-211
163	 <chem>CCN(Cc1ccc(cc1)C(=O)O)c2nc(c(s2)c3cc(Cl)ccc3OCc4ccccc4)cc5ccccc5</chem>	169-171

Table 40



Ex.	Structural formula	m.p. (°C)
164	 <chem>O=C(O)c1ccc(cc1)CN(C2CCCCC2)c3nc(cc3sc3c4ccc(Cl)cc4OCc5ccccc5)c6ccccc6</chem>	158- 160
165	 <chem>O=C(O)c1ccc(cc1)CN(C2CCCCC2)c3nc(cc3sc3c4ccc(Cl)cc4OCc5ccc(cc5)C(F)(F)F)c6ccccc6</chem>	171- 172

Table 41




Ex.	Structural formula	m.p. (°C)
166		183-184
167		196-198
168		143-149

Table 42

Ex.	Structural formula	m.p. (°C)
169	 <chem>CC1=CC=C(C=C1)OCC2=CC=C(C=C2)OC3=CC=C(C=C3)N(C3=CC=C(C=C3)C(=O)O)CC4CCCCC4</chem>	132– 137
170	 <chem>COc1cc(ccc1OC2=CC=C(C=C2)N(C2=CC=C(C=C2)C(=O)O)CC3CCCCC3)COC4=CC=C(C=C4)C(F)(F)F</chem>	104– 107

Table 43




Ex.	Structural formula	m.p. (°C)
171		111- 114
172		170- 171
173		215- 218

Table 44



Ex.	Structural formula	m.p. (°C)
174	 <chem>OC(=O)c1ccc(cc1)N(CC2CCCCC2)c3nc(cc3s3c4ccc(F)cc4OCC5=CC=CC=C5)cc3</chem>	125- 130
175	 <chem>OC(=O)c1ccc(cc1)N(CC2CCCCC2)c3nc(cc3s3c4ccc(C)cc4OCC5=CC=CC=C5)cc3</chem>	148- 150

Table 45

Ex.	Structural formula	m.p. (°C)
176	 <chem>CC(C)c1ccc(cc1)COc2cc(Cl)cc(cc2N3C(=N)SC(=N3)N(CCC4CCCCC4)c5ccc(cc5)C(=O)O)</chem>	190- 192
177	 <chem>FC(F)(F)c1ccc(cc1)COc2cc(Cl)cc(cc2N3C(=N)SC(=N3)N(CCC4CCCCC4)c5ccc(cc5)C(=O)O)</chem>	170- 175

Table 46



Ex.	Structural formula	m.p. (°C)
178		195- 198
179		134- 140

Table 47



Ex.	Structural formula	m.p. (°C)
180	 <chem>COc1cc(OCc2ccc(cc2)C(F)(F)F)cc1-c1cc[nH]c1-c1ccc(cc1)OC(=O)O</chem>	227-230
181	 <chem>Clc1cc(OCc2ccccc2)cc1-c1cc[nH]c1-c1ccc(cc1)CN(C1CCCCC1)C(=O)O</chem>	98-100

Table 48





Ex.	Structural formula	m.p. (°C)
182		216- 218
183		230
184		195- 197
185		173- 176

Table 49





Ex.	Structural formula	m.p. (°C)
186		220
187		182- 189
188		230 (dec.)
189		213- 214

Table 50





Ex.	Structural formula	m.p. (°C)
190		228-230
191		160-162
192		amorphous
193	 <p style="text-align: center;">ClH</p>	amorphous

Table 51

Ex.	Structural formula	m.p. (°C)
194		180-181
195		amorphous
196		194-196
197		amorphous
198		amorphous

Table 52





Ex.	Structural formula	m.p. (°C)
199		amorphous
200		amorphous
201		amorphous
202		amorphous

Table 53







Ex.	Structural formula	m.p. (°C)
203		amorphous
204		183 (dec.)
205		220
206		159-161
207		189-191
208		210-213

Table 54

Ex.	Structural formula	m.p. (°C)
209		190-193
210		225-227
211		208-210
212		181-183
213		191-193
214		amorphous

Table 55





Ex.	Structural formula	m.p. (°C)
215		155- 158
216		230
217		188- 190
218		185- 191

Table 56





Ex.	Structural formula	m.p. (°C)
219		amorphous
220		117-119
221		105-108
222		146-147

Table 57





Ex.	Structural formula	m.p. (°C)
223		174- 176
224		151- 152
225		127- 128
226		157- 159

Table 58





Ex.	Structural formula	m.p. (°C)
227		172- 173
228		155- 157
229		176- 177
230		114- 115

Table 59






Ex.	Structural formula	m.p. (°C)
231		250
232		144- 145
233		166- 167
234		180- 181
235		162

Table 60






Ex.	Structural formula	m.p. (°C)
236		158-159
237		164-167
238		212-213
239		159-163
240		130-135

Table 61





Ex.	Structural formula	m.p. (°C)
241		147- 148
242		149- 150
243		122- 127
244		153- 154

Table 62






Ex.	Structural formula	m.p. (°C)
245		132-134
246		205 (dec.)
247		209 (dec.)
248		104-105
249		117-118

Table 63






Ex.	Structural formula	m.p. (°C)
250		116-117
251		175 (dec.)
252		174 (dec.)
253		240
254		amorphous

Table 64






Ex.	Structural formula	m.p. (°C)
255		167- 170
256		168- 170
257		130 (dec.)
258		234- 236
259		231- 233

Table 65

Ex.	Structural formula	m.p. (°C)
260		220- 223
261		159 (dec.)
262		239 (dec.)
263		260
264		182- 183

Table 66

Ex.	Structural formula	m.p. (°C)
265		208- 221
266		200- 202
267		122- 123
268		88-90
269		99-101

Table 67

Ex.	Structural formula	m.p. (°C)
270		100-104
271		76-84
272		75-79
273		87-92
274		230 (dec.)

[0188] A Formulation Example is shown in the following, which is not to be construed as limitative.

Formulation Example

[0189]

(a) Compound of Example 1	10 g
(b) Lactose	50 g

(continued)

(c) Corn starch	15 g
(d) Sodium carboxymethyl cellulose	44 g
(e) Magnesium stearate	1 g

[0190] The total amount of (a), (b) and (c) and 30 g of (d) were kneaded with water, dried in vacuo and granulated. The granulated powder was mixed with 14 g of (d) and 1 g of (e) and the mixture was punched with a tableting machine to give 1000 tablets containing 10 mg of (a) per tablet.

[0191] The test results of the protein tyrosine phosphatase 1B inhibitory action of the present invention are shown in the following.

(Experimental Example)

Experimental Example 1 (protein tyrosine phosphatase 1B inhibitory action)

Preparation of assay buffer:

[0192] 50 mM Tris-HCl buffer (pH 7.5), and 50 mM NaCl and 3 mM dithiothreitol (DTT) were prepared.

Preparation of sample:

[0193] Respective 10 mM DMSO solutions of 0.1, 0.3, 1, 3 and 10 μ M test compounds were diluted with the above-mentioned assay buffer so that the final dimethyl sulfoxide (DMSO) concentration would be not more than 1%. As a control, an assay buffer was used.

Preparation of substrate:

[0194] A synthetic peptide consisting of 12 amino acids corresponding to 1142nd-1153rd of insulin receptor sequence, wherein three tyrosines therein had been phosphorylated, was diluted with the above-mentioned assay buffer to 80 μ M.

Preparation of enzyme:

[0195] A recombinant human protein tyrosine phosphatase 1B(manufactured by UBI) was diluted with the above-mentioned assay buffer (1.2 ng/25 μ l).

(Evaluation method)

[0196] A sample (10 μ l) and a substrate (25 μ l) prepared as mentioned above were successively added to a 96 well plate, and an enzyme (25 μ l) prepared as mentioned above was added and mixed. After incubation at room temperature for 60 min., malachite green (120 μ l, Biomol), which is a phosphorus color fixing agent, was added, and the mixture was further incubated at room temperature for 20 min. to allow color development. The absorbance at 650 nm was measured on a plate reader, and the protein tyrosine phosphatase 1B inhibitory action of the test compound was evaluated. The results are shown in Table 68.

Experimental Example 2 (hypoglycemic action)

[0197] A 0.5% methyl cellulose suspension of the test compound was orally administered to male ob/ob mice (6 to 9 weeks old) grouped based on blood glucose level. A 0.5% methyl cellulose solution alone was administered to a control group.

[0198] The blood was drawn from the orbital vein with anesthesia at 3 hours after the administration. The blood was drawn under fasting by removing the feed immediately before test compound administration. The blood thus drawn was separated by centrifugation and the blood glucose level was measured from the obtained plasma according to the hexokinase method (glucose measure kit). For evaluation, decrease in the blood glucose level of the test compound administration group relative to the control group was shown in percentage. The results are shown in Table 68.

Table 68

Example	PTP1B inhibitory action (IC ₅₀ : μ M)	Decrease in blood glucose level (%)	
		Dose (mg/kg)	3 hr
1	0.32	0.3	35
		1	35
		3	35
2	0.22	1	37
		3	47
3	0.50	1	27
		3	35
115	0.48	3	34
196	0.83	3	-8
208	0.11	3	21
209	0.49	3	20
215	0.11	0.3	-9
		1	32
		3	39
217	0.26	3	20
218	0.15	3	27
230	0.15	3	40

Experimental Example 3 (blood lipid lowering action)

[0199] A 0.5% methyl cellulose suspension of the test compound was orally administered to 7-week-old db/db mice once a day for 14 days. The 0.5% methyl cellulose solution alone was administered to the control group.

[0200] The blood was drawn from the orbital vein with light ether anesthesia before test compound administration on day 7 and day 14 under non-fasting condition. The blood thus drawn was separated by centrifugation and the plasma triglyceride concentration was measured from the obtained plasma according to the enzyme method (triglyceride measure kit). The results are shown in Table 69.

Table 69

	Dose (mg/kg)	Plasma triglyceride concentration (mg/dL)	
		Day 7	Day 14
Control group	-----	508	622
test compound administration group (Example 102)	10	277	349
	30	250	307


Industrial Applicability

[0201] From the foregoing test results and the like, the compound [I] of the present invention is suggested have a superior PTP1B inhibitory action. That is, the compound [I] of the present invention is expected to be a new type of drug for the prophylaxis or treatment of diabetes that can directly improve the insulin action, insulin sensitivity, insulin resistance and/or glucose resistance. In addition, the compound [I] of the present invention is also expected as a drug for the prophylaxis or treatment of diabetic complications (retinopathy, nephropathy, neuropathy, cardiac infarction and cerebral infarction based on arteriosclerosis etc.), and further as a treatment drug of a disease mediated by PTP1B. Moreover, because compound [I] of the present invention has been confirmed to show a blood lipid lowering action from the above test results, it is also expected as a drug for the prophylaxis or treatment of hyperlipidemia.

[0202] This application is based on a patent application No. 368567/2001 filed in Japan, the contents of which are hereby incorporated by reference.

Claims

1. An azole compound represented by the formula [I]



wherein

W is a sulfur atom or an oxygen atom;

R is

(1) $-\text{COOR}^7$ wherein R^7 is a hydrogen atom or a lower alkyl group),

(2) $-\text{X}^1-\text{A}^1-\text{COOR}^7$

wherein

X^1 is $-\text{O}-$, $-\text{N}(\text{R}^{15})-$ or $-\text{S}(\text{O})_p-$ wherein R^{15} is a hydrogen atom or a lower alkyl group, p is 0, 1 or 2,

A^1 is a lower alkylene group, and

R^7 is a hydrogen atom or a lower alkyl group, or

(3) a tetrazolyl group;

R^1 , R^2 , R^3 and R^4 are each independently

(1) a hydrogen atom,

(2) a halogen atom,

(3) a hydroxyl group,

(4) an optionally substituted lower cycloalkylalkyloxy group,

(5) an optionally substituted aralkyloxy group,

(6) a cyano group,

(7) a nitro group,

(8) a lower alkyl group,

(9) a lower haloalkyl group,

(10) a lower alkoxy group or

(11) a lower haloalkoxy group;

A is a group represented by $-(\text{CH}_2)_m-\text{X}-$

wherein

X is $-\text{N}(\text{R}^8)-$, $-\text{C}(\text{R}^9)(\text{R}^{10})-$, $-\text{CO}-$ or $-\text{CO}-\text{N}(\text{R}^8)-$

wherein

R^8 is a hydrogen atom, $-\text{SO}_2\text{R}^{16}$ (R^{16} is a lower alkyl group or an aryl group) or a lower alkyl group, wherein said lower alkyl group is optionally substituted by a substituent selected from the group consisting of a lower alkoxy group, an aryloxy group, $-\text{N}(\text{R}^{11})(\text{R}^{12})$ (R^{11} and R^{12} are each independently a hydrogen atom or a lower alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a lower cycloalkyl group and an optionally substituted aryl group, and

R^9 and R^{10} are each independently a hydrogen atom or a lower alkyl group or may form lower cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, and

m is 0 or an integer of 1 to 3;

B is an aryl group or an aromatic heterocyclic group;

R^5 is

(1) a hydrogen atom,

- (2) a halogen atom,
 (3) a lower alkyl group,
 (4) a lower alkoxy group,
 (5) a cyano group,
 (6) a nitro group,
 (7) a lower haloalkyl group or
 (8) $-S(O)_r-R^{17}$ (R^{17} is a lower alkyl group or an aryl group and r is 0, 1 or 2); and

R^6 is $-(Y)_{s1}-(A^2)_s-Z$

wherein

$s1$ and s are each independently 0 or 1,

Y is $-O-$, $-S(O)_t-$, $-N(R^{13})-$, $-N(R^{14})-CO-$, $-N(R^{14})-SO_2-$, $-SO_2-N(R^{14})-$, $-C(R^{18})(R^{19})-$ or $-CO-$
 (wherein

t is 0, 1 or 2,

R^{13} is

- (1) a hydrogen atom,
 (2) a lower alkyl group
 (wherein said lower alkyl group optionally substituted by a substituent selected from the group consisting of

- (a) a lower cycloalkyl group,
 (b) an optionally substituted aryl group,
 (c) an optionally substituted heterocyclic group and
 (d) a hydroxyl group),

- (3) a lower alkenyl group,
 (4) a lower alkylsulfonyl group or
 (5) a lower alkylcarbonyl group

(wherein said lower alkylcarbonyl group is optionally substituted by a hydroxyl group or a lower alkoxy group),

R^{14} is a hydrogen atom or a lower alkyl group, and

R^{18} and R^{19} are each independently a hydrogen atom or a lower alkyl group or may form lower cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom),

A^2 is a lower alkylene group optionally substituted by a lower cycloalkyl group, and

Z is

- (1) a lower cycloalkyl group
 (wherein said a lower cycloalkyl group is optionally substituted by an optionally substituted phenyl group),
 (2) an aryl group
 (wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group optionally substituted by a substituent selected from the group consisting of a lower alkyl group and a lower alkylcarbonyl group,
 (b) a lower cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a lower alkyl group,
 (c) a carboxy group,
 (d) a halogen atom,
 (e) an alkyl group,
 (f) a lower haloalkyl group,
 (g) a lower alkylamino group,
 (h) a di(lower alkyl)amino group,
 (i) a lower alkylthio group and
 (j) a lower alkoxy group),

- (3) an optionally substituted aromatic heterocyclic group,
 (4) an indanyl group or

(5) a piperazinyl group

(wherein said piperazinyl group is optionally substituted by a substituent selected from the group consisting of

- 5 (a) a phenyl group,
 (b) a phenyl lower alkyl group,
 (c) a benzoyl group optionally substituted by a halogen atom and
 (d) a phenyl lower alkoxy carbonyl group),

10 a prodrug thereof or a pharmaceutically acceptable salt thereof.

2. The azole compound of claim 1, wherein, in the formula [I], W is a sulfur atom or an oxygen atom;
 R is

- 15 (1) $-\text{COOR}^7$ wherein R^7 is a hydrogen atom or a C_{1-4} alkyl group,
 (2) $-\text{X}^1\text{-A}^1\text{-COOR}^7$
 wherein
 X^1 is $-\text{O}-$, $-\text{N}(\text{R}^{15})-$ or $-\text{S}(\text{O})_p-$ wherein R^{15} is a hydrogen atom or a C_{1-4} alkyl group and p is 0, 1 or 2,
 A^1 is a C_{1-4} alkylene group,
 20 R^7 is a hydrogen atom or a C_{1-4} alkyl group or
 (3) a tetrazolyl group,

R^1 , R^2 , R^3 and R^4 are each independently,

- 25 (1) a hydrogen atom,
 (2) a halogen atom,
 (3) a hydroxyl group,
 (4) an optionally substituted C_{3-7} cycloalkyl C_{1-4} alkoxy group,
 (5) an optionally substituted aralkoxy group,
 30 (6) a cyano group,
 (7) a nitro group,
 (8) a C_{1-4} alkyl group,
 (9) a C_{1-4} haloalkyl group,
 (10) a C_{1-4} alkoxy group or
 35 (11) a C_{1-4} haloalkoxy group;

A is a group represented by $-(\text{CH}_2)_m\text{-X-}$
 wherein

X is $-\text{N}(\text{R}^8)-$, $-\text{C}(\text{R}^9)(\text{R}^{10})-$, $-\text{CO-}$ or $-\text{CO-N}(\text{R}^8)-$

40 wherein

R^8 is a hydrogen atom, $-\text{SO}_2\text{R}^{16}$ (R^{16} is a C_{1-6} alkyl group or an aryl group) or a C_{1-6} alkyl group, wherein said C_{1-6} alkyl group is optionally substituted by a substituent selected from the group consisting of a C_{1-4} alkoxy group, an aryloxy group, $-\text{N}(\text{R}^{11})(\text{R}^{12})$ (R^{11} and R^{12} are each independently a hydrogen atom or a C_{1-4} alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, C_{3-7} cycloalkyl group and an optionally substituted aryl group,

45 R^9 and R^{10} are each independently a hydrogen atom or a C_{1-4} alkyl group, or may form a C_{3-7} cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, m is 0 or an integer of 1 to 3;

B is an aryl group or an aromatic heterocyclic group;

R^5 is

- 55 (1) a hydrogen atom,
 (2) a halogen atom,
 (3) a C_{1-4} alkyl group,
 (4) a C_{1-4} alkoxy group,
 (5) a cyano group,

- (6) a nitro group,
 (7) a C₁₋₄ haloalkyl group or
 (8) -S(O)_r-R¹⁷ (R¹⁷ is a C₁₋₆ alkyl group or an aryl group and r is 0, 1 or 2);

5 R⁶ is -(Y)_{s1}-(A²)_s-Z

wherein

s₁ and s are each independently 0 or 1,

Y is -O-, -S(O)_t-, -N(R¹³)-, -N(R¹⁴)-CO-, -N(R¹⁴)-SO₂-, -SO₂-N(R¹⁴)-, -C(R¹⁸)(R¹⁹)- or -CO-

(wherein

10 t is 0, 1 or 2,

R¹³ is

(1) a hydrogen atom,

(2) a C₁₋₄ alkyl group

15 (wherein said C₁₋₄ alkyl group is optionally substituted by a substituent selected from the group consisting of

(a) a C₃₋₇ cycloalkyl group,

(b) an optionally substituted aryl group,

(c) an optionally substituted heterocyclic group and

20 (d) a hydroxyl group),

(3) a C₂₋₄ alkenyl group,

(4) a C₁₋₄ alkylsulfonyl group or

(5) a C₁₋₄ alkylcarbonyl group

25 (wherein said C₁₋₄ alkylcarbonyl group is optionally substituted by a hydroxyl group or a C₁₋₄ alkoxy group),

R¹⁴ is a hydrogen atom or a C₁₋₄ alkyl group,

30 R¹⁸ and R¹⁹ are each independently a hydrogen atom or a C₁₋₄ alkyl group, or may form C₃₋₇ cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom),

A² is a C₁₋₄ alkylene group optionally substituted by a C₃₋₇ cycloalkyl group,

Z is

35 (1) a C₃₋₇ cycloalkyl group

(wherein said C₃₋₇ cycloalkyl group is optionally substituted by a phenyl group optionally substituted by a halogen atom),

(2) an aryl group

(wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

40 (a) a heterocyclic group optionally substituted by a substituent selected from the group consisting of a C₁₋₄ alkyl group and a C₁₋₄ alkylcarbonyl group,

(b) a C₃₋₇ cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,

45 (c) a carboxy group,

(d) a halogen atom,

(e) a C₁₋₈ alkyl group,

(f) a C₁₋₄ haloalkyl group,

(g) a C₁₋₄ alkylamino group,

50 (h) a di(C₁₋₄ alkyl)amino group,

(i) a C₁₋₄ alkylthio group and

(j) a C₁₋₄ alkoxy group),

(3) an aromatic heterocyclic group

55 (wherein said aromatic heterocyclic group is optionally substituted by a substituent selected from the group consisting of

(a) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group,

- (b) a C₁₋₆ alkyl group,
- (c) an aryl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group,
- (d) a halogen atom,
- (e) a C₁₋₄ haloalkyl group,
- (f) a carboxy group,
- (g) a C₃₋₇ cycloalkyl group and
- (h) a C₁₋₄ alkoxy group),

- (4) an indanyl group or
- (5) a piperazinyl group

(wherein said piperazinyl group is optionally substituted by a substituent selected from the group consisting of

- (a) a phenyl group,
- (b) a phenyl C₁₋₄ alkyl group,
- (c) a benzoyl group optionally substituted by a halogen atom and
- (d) a phenyl C₁₋₄ alkoxy carbonyl group),

a prodrug thereof or a pharmaceutically acceptable salt thereof.

3. The azole compound of claim 2, wherein W is a sulfur atom or an oxygen atom,
R is

- (1) -COOR⁷ wherein R⁷ is a hydrogen atom,
 - (2) -X¹-A¹-COOR⁷
- wherein
X¹ is -O-,
A¹ is a C₁₋₄ alkylene group,
R⁷ is a hydrogen atom or
- (3) a tetrazolyl group;

R¹, R², R³ and R⁴ are each independently

- (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a hydroxyl group,
- (4) an optionally substituted C₃₋₇ cycloalkyl C₁₋₄ alkyloxy group or
- (5) an optionally substituted aralkyloxy group;

A is a group represented by -(CH₂)_m-X-
wherein
X is -N(R⁸)-, -C(R⁹)(R¹⁰)- or -CO-
wherein

R⁸ is a hydrogen atom or a C₁₋₆ alkyl group, wherein said C₁₋₆ alkyl group is optionally substituted by a substituent selected from the group consisting of a C₁₋₄ alkoxy group, an aryloxy group, -N(R¹¹)(R¹²) (R¹¹ and R¹² are each independently a hydrogen atom or a C₁₋₄ alkyl group or may form, together with a nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a C₃₋₇ cycloalkyl group and an optionally substituted aryl group,

R⁹ and R¹⁰ are each independently a hydrogen atom or a C₁₋₄ alkyl group, or may form C₃₋₇ cycloalkane together with a carbon atom bonded thereto, or may form, together with a carbon atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, m is 0 or an integer of 1 to 3;

B is an aryl group or an aromatic heterocyclic group;

R⁵ is

- (1) a hydrogen atom,
- (2) a halogen atom,

- (3) a C₁₋₄ alkyl group or
- (4) a C₁₋₄ alkoxy group;

R⁶ is -(Y)_{s1}-(A²)_s-Z

wherein

s₁ and s are each independently 0 or 1,

Y is -O-, -S(O)_t-, -N (R¹³)-, -N (R¹⁴)-CO- or -N(R¹⁴)-SO₂-

wherein

t is 0, 1 or 2,

R¹³ is

- (1) a hydrogen atom,
- (2) a C₁₋₄ alkyl group
- (wherein said C₁₋₄ alkyl group is optionally substituted by a substituent selected from the group consisting of

- (a) a C₃₋₇ cycloalkyl group,
- (b) an optionally substituted aryl group,
- (c) an optionally substituted heterocyclic group and
- (d) a hydroxyl group),

- (3) a C₂₋₄ alkenyl group,
- (4) a C₁₋₄ alkylsulfonyl group or
- (5) a C₁₋₄ alkylcarbonyl group

(wherein said C₁₋₄ alkylcarbonyl group is optionally substituted by a hydroxyl group or a C₁₋₄ alkoxy group),

R¹⁴ is a hydrogen atom or a C₁₋₄ alkyl group,

A² is a C₁₋₄ alkylene group optionally substituted by a C₃₋₇ cycloalkyl group,

Z is

- (1) a C₃₋₇ cycloalkyl group
- (wherein said C₃₋₇ cycloalkyl group is optionally substituted by a phenyl group),
- (2) an aryl group
- (wherein said aryl group is optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group or a C₁₋₄ alkylcarbonyl group,
- (b) a C₃₋₇ cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,
- (c) a carboxy group,
- (d) a halogen atom,
- (e) a C₁₋₈ alkyl group,
- (f) a C₁₋₄ haloalkyl group,
- (g) a C₁₋₄ alkylamino group and
- (h) a di (C₁₋₄ alkyl) amino group,
- (i) a C₁₋₄ alkylthio group and
- (j) a C₁₋₄ alkoxy group),

- (3) an aromatic heterocyclic group
- (wherein said aromatic heterocyclic group is optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group,
- (b) a C₁₋₄ alkyl group and
- (c) a phenyl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group),

- (4) an indanyl group or
- (5) a piperazinyl group

(wherein said piperazinyl group is optionally substituted by a substituent selected from the group consisting of

- (a) a phenyl group,
- (b) a phenyl C₁₋₄ alkyl group and
- (c) a phenyl C₁₋₄ alkoxy carbonyl group),

- 5 a prodrug thereof or a pharmaceutically acceptable salt thereof.
4. The azole compound of claim 3, wherein W is a sulfur atom and m is 0 or 1, a prodrug thereof or a pharmaceutically acceptable salt thereof.
- 10 5. The azole compound of claim 4, wherein A is -(CH₂)_m-X-
wherein
X is -N(R⁸)- wherein R⁸ is a hydrogen atom or a C₁₋₆ alkyl group, wherein said C₁₋₆ alkyl group is optionally substituted by a substituent selected from the group consisting of a C₁₋₄ alkoxy group, an aryloxy group, -N(R¹¹) (R¹²) (R¹¹ and R¹² are each independently a hydrogen atom or a C₁₋₄ alkyl group or may form, together with a
15 nitrogen atom bonded thereto, a 5- to 7-membered hetero ring optionally further having at least one hetero atom selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), a carboxy group, a C₃₋₇ cycloalkyl group and an optionally substituted aryl group, and m is 0 or 1,
a prodrug thereof or a pharmaceutically acceptable salt thereof.
- 20 6. The azole compound of claim 5, wherein R is -X¹-A¹-COOR⁷ wherein each symbol is as defined in claim 3,
a prodrug thereof or a pharmaceutically acceptable salt thereof.
7. The azole compound of claim 5, wherein R is -COOR⁷ wherein R⁷ is a hydrogen atom,
a prodrug thereof or a pharmaceutically acceptable salt thereof.
- 25 8. The azole compound of claim 7, wherein R¹, R², R³ and R⁴ are hydrogen atoms,
a prodrug thereof or a pharmaceutically acceptable salt thereof.
9. The azole compound of claim 8, wherein B is a phenyl group, a thiazolyl group, a pyridyl group, a benzothiazolyl group, a benzoimidazolyl group or a benzoxazolyl group,
30 a prodrug thereof or a pharmaceutically acceptable salt thereof.
10. The azole compound of claim 9, wherein B is a phenyl group, a prodrug thereof or a pharmaceutically acceptable salt thereof.
- 35 11. The azole compound of claim 10, wherein R⁵ is a hydrogen atom,
a prodrug thereof or a pharmaceutically acceptable salt thereof.
- 40 12. The azole compound of claim 11, wherein, for R⁶, Z is
- (1) a C₃₋₇ cycloalkyl group
(wherein said C₃₋₇ cycloalkyl group is optionally substituted by a phenyl group) ,
 - (2) an aryl group
(wherein said aryl group is optionally substituted by a substituent selected from the group consisting of
 - 45 (a) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group or a C₁₋₄ alkyl carbonyl group,
 - (b) a C₃₋₇ cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,
 - (c) a carboxy group,
 - 50 (d) a halogen atom,
 - (e) a C₁₋₈ alkyl group,
 - (f) a C₁₋₄ haloalkyl group,
 - (g) a C₁₋₄ alkylamino group,
 - (h) a di(C₁₋₄ alkyl)amino group,
 - 55 (i) a C₁₋₄ alkylthio group and
 - (j) a C₁₋₄ alkoxy group) or
 - (3) an aromatic heterocyclic group

(wherein said aromatic heterocyclic group is optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group,
- (b) a C₁₋₄ alkyl group and
- (c) a phenyl group optionally substituted by a halogen atom or a C₁₋₄ haloalkyl group),

a prodrug thereof or a pharmaceutically acceptable salt thereof.

13. The azole compound of claim 12, wherein Z is an aryl group optionally substituted by a substituent selected from the group consisting of

- (a) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group or a C₁₋₄ alkylcarbonyl group,
- (b) a C₃₋₇ cycloalkyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,
- (c) a carboxy group,
- (d) a halogen atom,
- (e) a C₁₋₈ alkyl group,
- (f) a C₁₋₄ haloalkyl group,
- (g) a C₁₋₄ alkylamino group,
- (h) a di(C₁₋₄ alkyl)amino group,
- (i) a C₁₋₄ alkylthio group and
- (j) a C₁₋₄ alkoxy group,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

14. The azole compound of claim 13, wherein Z is a phenyl group substituted by a substituent selected from the group consisting of

- (a) a cyclohexyl group or a cyclopentyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,
 - (b) a heterocyclic group optionally substituted by a C₁₋₄ alkyl group or a C₁₋₄ alkylcarbonyl group (wherein said heterocyclic group is selected from the group consisting of a piperidiny group, a morpholinyl group, a piperazinyl group, a tetrahydropyranyl group, a pyrrolidinyl group and a pyrrolyl group) and
 - (c) a C₁₋₈ alkyl group,
- a prodrug thereof or a pharmaceutically acceptable salt thereof.

15. The azole compound of claim 14, wherein Z is a phenyl group substituted by a cyclohexyl group optionally substituted by a substituent selected from the group consisting of a hydroxyl group, an oxo group, a halogen atom and a C₁₋₄ alkyl group,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

16. The azole compound of claim 13 or claim 14 wherein, for R⁶, Y is -O-, -N(R¹³)- or -N(R¹⁴)-CO- wherein

R¹³ is a hydrogen atom, a C₁₋₄ alkyl group or a C₂₋₄ alkenyl group, wherein said C₁₋₄ alkyl group is optionally substituted by a substituent selected from the group consisting of a C₃₋₇ cycloalkyl group, an optionally substituted aryl group, an optionally substituted heterocyclic group and a hydroxyl group, R¹⁴ is a hydrogen atom or a C₁₋₄ alkyl group, and s1 is 1,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

17. The azole compound of claim 16, wherein, for R⁶, A² is a methylene group,

a prodrug thereof or a pharmaceutically acceptable salt thereof.

18. A pharmaceutical composition comprising an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

19. A pharmaceutical composition for inhibiting protein Tyrosine Phosphatase 1B, which comprises an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically

acceptable carrier.

20. A pharmaceutical composition for treating diabetes, which comprises an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
21. A pharmaceutical composition for treating hyperlipidemia, which comprises an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
22. The pharmaceutical composition of claim 18, which is used in combination with a different therapeutic drug for hyperlipidemia.
23. The pharmaceutical composition of claim 22, wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.
24. The pharmaceutical composition of claim 23, wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.
25. The pharmaceutical composition of claim 18, which is used in combination with a different therapeutic drug for diabetes.
26. The pharmaceutical composition of claim 25, which is used in combination with a therapeutic agent for diabetes selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin preparation and an insulin sensitizer.
27. The pharmaceutical composition of claim 26, wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glycopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochloride.
28. The pharmaceutical composition of claim 20, which is used in combination with a different therapeutic drug for hyperlipidemia.
29. The pharmaceutical composition of claim 28, wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.
30. The pharmaceutical composition of claim 29, wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.
31. The pharmaceutical composition of claim 20, which is used in combination with a different therapeutic drug for diabetes.
32. The pharmaceutical composition of claim 31, which is used in combination with a therapeutic drug for diabetes selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin preparation and an insulin sensitizer.
33. The pharmaceutical composition of claim 32, wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glycopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochloride.
34. The pharmaceutical composition of claim 21, which is used in combination with a different therapeutic drug for hyperlipidemia.
35. The pharmaceutical composition of claim 34, wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.
36. The pharmaceutical composition of claim 35, wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.

37. The pharmaceutical composition of claim 21, which is used in combination with a different therapeutic drug for diabetes.
- 5 38. The pharmaceutical composition of claim 37, which is used in combination with a therapeutic drug for diabetes selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin preparation and an insulin sensitizer.
- 10 39. The pharmaceutical composition of claim 38, wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glycopyramide, tolazamide, glybuzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochloride.
- 15 40. A method of inhibiting protein Tyrosine Phosphatase 1B, which comprises administering an effective amount of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal.
41. A method of treating diabetes, which comprises administering an effective amount of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal.
- 20 42. A method of treating hyperlipidemia, which comprises administering an effective amount of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal.
43. Use of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof for the production of a protein tyrosine phosphatase 1B inhibitor.
- 25 44. Use of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof for the production of a therapeutic agent for diabetes.
45. Use of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof for the production of a therapeutic agent for hyperlipidemia.
- 30 46. A commercial package comprising the pharmaceutical composition of claim 18 and a written matter associated therewith, the written matter stating that said pharmaceutical composition can or should be used for inhibiting protein Tyrosine Phosphatase 1B.
- 35 47. A commercial package comprising the pharmaceutical composition of claim 18 and a written matter associated therewith, the written matter stating that said pharmaceutical composition can or should be used for treating diabetes.
- 40 48. A commercial package comprising the pharmaceutical composition of claim 18 and a written matter associated therewith, the written matter stating that said pharmaceutical composition can or should be used for treating hyperlipidemia.
- 45 49. A method of treating hyperlipidemia, which comprises administering an effective amount of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal, and administering an effective amount of a different therapeutic drug for hyperlipidemia to said mammal.
50. The method of treating of claim 49, wherein the therapeutic drug for hyperlipidemia is a statin pharmaceutical agent.
- 50 51. The method of treating of claim 50, wherein the statin pharmaceutical agent is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin.
52. A method of treating diabetes, which comprises administering an effective amount of an azole compound of any of claims 1 to 17, a prodrug thereof or a pharmaceutically acceptable salt thereof to a mammal, and administering an effective amount of a different therapeutic agent for diabetes to said mammal.
- 55 53. The method of treating of claim 52, wherein the therapeutic agent for diabetes is selected from the group consisting of an insulin secretagogue, a sulfonylurea, a sulfonamide, a biguanide, an α glucosidase inhibitor, an insulin prep-

aration and an insulin sensitizer.

54. The method of treating of claim 53, wherein the therapeutic agent for diabetes is selected from the group consisting of nateglide, glimepiride, glibenclamide, gliclazide, acetohexamide, tolbutamide, glycopyramide, tolazamide, gly-
5 buzole, metformin hydrochloride, buformin hydrochloride, voglibose, acarbose, insulin and pioglitazone hydrochlo-
ride.

10

15

20

25

30

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12673

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl⁷ C07D277/42, 417/04, 417/12, A61K31/426, 31/427, 31/454,
31/496, 31/5377, 31/4439, A61P3/06, 3/10, 43/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl⁷ C07D277/42, 417/04, 417/12, A61K31/426, 31/427, 31/454,
31/496, 31/5377, 31/4439, A61P3/06, 3/10, 43/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAPLUS (STN), CAOLD (STN), REGISTRY (STN), WPI/L (DIALOG)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6011048 A (MERCK AND CO., INC.), 04 January, 2001 (04.01.00), Claims; column 20, line 57 to column 21, line 15; column 23, lines 8 to 62	1-2, 18, 20-39, 44-45, 47-48
Y	& EP 968209 A1 & WO 98/32753 A1 & JP 2001-509166 A	18-39, 43-48
Y	JP 2000-504039 A (Sanofi), 04 April, 2000 (04.04.00), Claims; page 28, line 18 to page 29, line 4 & WO 98/15543 A1 & EP 934290 A1 & US 6344470 A	1-5, 7-13, 18-39, 43-48
Y	WO 99/58511 A1 (AMERICAN HOME PRODUCTS CORP.), 18 November, 1999 (18.11.99), Claims; page 11, line 9 to page 17, line 8 & EP 1077958 A1 & JP 2002-505317 A	1-5, 7-13, 18-39, 43-48

☒ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not
considered to be of particular relevance

"E" earlier document but published on or after the international filing
date

"L" document which may throw doubts on priority claim(s) or which is
cited to establish the publication date of another citation or other
special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other
means

"P" document published prior to the international filing date but later
than the priority date claimed

"T" later document published after the international filing date or
priority date and not in conflict with the application but cited to
understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be
considered novel or cannot be considered to involve an inventive
step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be
considered to involve an inventive step when the document is
combined with one or more other such documents, such
combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
07 February, 2003 (07.02.03)

Date of mailing of the international search report
04 March, 2003 (04.03.03)

Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12673

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5643932 A (OTSUKA PHARMACEUTICAL CO., LTD.), 01 July, 1997 (01.07.97), Claims; column 3, line 61 to column 4, line 17 & EP 513387 A1 & JP 5-51318 A	1-5, 7-13, 18-39, 43-48

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12673

Box I Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 40 to 42, 49 to 54

because they relate to subject matter not required to be searched by this Authority, namely:

Claims 40 to 42, 49 to 54 pertain to a method for treatment of the human body by therapy and thus relate to a subject matter which this International Searching Authority is not required, under the provisions of Article 17(2)(a)(i) of (continued to extra sheet)

2. ☐ Claims Nos.:

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:

because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12673

Continuation of Box No.I-1 of continuation of first sheet(1)

the PCT and Rule 39.1(iv) of the Regulations under the PCT, to make an international search.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.